

Ancient Solutions to Geometric Flows

Panagiota Daskalopoulos

Columbia University

Recent trends in Nonlinear Evolution Equations

CIRM

April 4-8, 2016

Ancient and Eternal Solutions

- We will discuss **ancient** or **eternal** solutions to **geometric parabolic** partial differential equations.
- These are **special** solutions which exist for all time

$$-\infty < t < T \quad \text{where } T \in (-\infty, +\infty].$$

- They appear as **blow up** limits near a **singularity**.
- Understanding ancient and eternal solutions often sheds new insight to the **singularity analysis**

In this talk we will address:


- the **classification** of **ancient** solutions to **parabolic** partial differential equations, with emphasis to **geometric flows**: **Mean Curvature** flow, **Ricci** flow and **Yamabe** flow.
- methods of **constructing** new ancient solutions from the **gluing** of two or more **solitons** (self-similar solutions).
- new techniques and future research directions.

Ancient and Eternal solutions


- **Definition:** A solution $u(\cdot, t)$ to a parabolic equation is called **ancient** if it is defined for all time $-\infty < t < T$, $T < +\infty$.
- **Ancient** solutions typically arise as blow up limits at a **type I** singularity.
- **Definition:** A solution $u(\cdot, t)$ to a parabolic equation is called **eternal** if it is defined for all $-\infty < t < +\infty$.
- **Eternal** solutions as blow up limits at a **type II** singularity.

Solitons



- **Solitons** (self-similar solutions) are typical examples of ancient or eternal solutions and often **models of singularities**.
- Some typical **examples** of **solitons** to geometric PDE are:
- **Spheres:**



- **Cylinders:**



- **Translating solitons:**




Solitons as singularities

- A well known technique introduced by R. Hamilton (1995) has been widely used to characterize as solitons the eternal solutions to geometric flows which attain a space-time curvature maximum.
- Such solutions typically appear as *carefully chosen blow up limits* near type II singularities.
- Its proof relies on a clever combination of the strong maximum principle and Li-Yau type differentiable Harnack estimates.

Other Ancient and eternal solutions

- However, there exist other ancient or eternal solutions which are **not solitons**.
- These, often may be visualized as obtained from the **gluing** as $t \rightarrow -\infty$ of two or more solitons.



- Obtaining more information about such solutions, often leads to the better understanding of the singularities.
- **Objective:** How to **construct** such solutions and how to **characterize** them among all ancient or eternal solutions.

Geometric conditions of ancient or eternal solutions

Goal: Characterize all **ancient** or **eternal** solutions to a **geometric flow** under natural **geometric conditions**:

- Being a **soliton** (self-similar solution)
- Satisfying an appropriate **curvature bound** as $t \rightarrow -\infty$:
 - i. **Type I**: **global curvature bound** after **typical** scaling.
 - ii. **Type II**: solutions which are **not type I**.
- Satisfying a **non-collapsing** condition.

The Semi-linear heat equation

- Consider next the **semilinear heat equation**

$$(\star_{SL}) \quad u_t = \Delta u + u^p \quad \text{on } \mathbb{R}^n \times (0, T)$$

in the **subcritical** range of exponents $1 < p < \frac{n+2}{n-2}$.

- It provides a prototype for the **blow up** analysis of **geometric flows**.
- In particular in **neckpinches** of solutions to the **Ricci flow** and **Mean Curvature flow**.
- Also in the characterization of **rescaled limits** as $t \rightarrow -\infty$ of **ancient solutions**.

The rescaled semi-linear heat equation

- Self-similar scaling at a singularity at (a, T) :

$$w(y, \tau) = (T-t)^{\frac{1}{p-1}} u(x, t), \quad y = \frac{x-a}{\sqrt{T-t}}, \quad \tau = -\log(T-t).$$

- Giga - Kohn (1985): $\|w(\tau)\|_{L^\infty(\mathbb{R}^n)} \leq C, \tau > -\log T.$
- The rescaled solution satisfies the equation

$$(\star) \quad w_\tau = \Delta w - \frac{1}{2}y \cdot \nabla w - \frac{w}{p-1} + w^p.$$

- **Objective:** To analyze the blow up behavior of u one needs to understand the long time behavior of w as $\tau \rightarrow +\infty.$
- This is closely related to the classification of bounded eternal solutions of $(\star).$

Eternal solutions of the semi-linear heat equation

- **Problem:** Provide the classification of **bounded** positive **eternal** solutions w of equation

$$(\star) \quad w_\tau = \Delta w - \frac{1}{2}y \cdot \nabla w - \frac{w}{p-1} + w^p.$$

- **Eternal** means that $w(\cdot, \tau)$ is defined for $\tau \in (-\infty, +\infty)$.
- The only **steady states** of (\star) are the **constants**:

$$w = 0 \quad \text{or} \quad w = \kappa, \quad \text{with} \quad \kappa := (p-1)^{-\frac{1}{(p-1)}}.$$

- **Theorem** (Giga - Kohn '87) $\lim_{\tau \rightarrow \pm\infty} w(\cdot, \tau) = \text{steady state}.$
- **Space independent eternal solutions** : $\phi(\tau) = \kappa(1 + e^\tau)^{-\frac{1}{(p-1)}}.$

Classification of Eternal solutions

- **Theorem** (Giga - Kohn '87 and Merle - Zaag '98)

If w is bounded positive **eternal** solution of (\star) defined on $\mathbb{R}^n \times (-\infty, +\infty)$, then

$$w = 0 \text{ or } w = \kappa \text{ or } w = \phi(\tau - \tau_0).$$

- **Main difficulty** (Merle - Zaag): Classify the orbits $w(\cdot, \tau)$ that connect the two **steady states**:

$$\lim_{\tau \rightarrow -\infty} w(\cdot, \tau) = \kappa \quad \text{and} \quad \lim_{\tau \rightarrow +\infty} w(\cdot, \tau) = 0$$

- **Elliptic analogues**: In similar spirit as the well known results by Gidas-Ni-Nirenberg '79 and Berestycki-Nirenberg '91 for global solutions to the elliptic equation

$$\Delta u + u^p = 0, \quad \text{on } \mathbb{R}^n.$$

- Other **Liouville type** results related to equation (\star_{SL}) by: P. Polacik, P. Quittner, T. Bartsch, P. Souplet, E. Yanagida among others.

Liouville type results for solutions to parabolic equations

- Although **Liouville type results** often appear with respect to **elliptic equations**, there are not many such results available in the **parabolic setting**.
- G. Koch, N. Nadirashvili, G. Seregin and V. Sverak (2009):
 - (i) **Liouville type** result for ancient **bounded** solutions of the **2-dim Navier Stokes** equations.
 - (ii) Also, similar result for bounded, **axi-symmetric with no swirl** solutions of the **3-dim Navier Stokes** equations.
- The proof relies on a weak type variation of the strong maximum principle for weak solutions of

$$u_t + a(x, t) \cdot \nabla u - \Delta u = 0.$$

- This result is applied on the vorticity $\omega := \operatorname{curl} u$.

Ancient Convex solutions to the CSF

- Let Γ_t be a family of closed curves which is an **embedded** solution to the **Curve shortening flow**, i.e. the embedding $F : \Gamma_t \rightarrow \mathbb{R}^2$ satisfies

$$\frac{\partial F}{\partial t} = -\kappa \nu$$

with κ the **curvature** of the curve and ν the **outer normal**.



- Gage (1984); Gage and Hamilton (1996); Grayson (1987):** the CSF shrinks Γ_t to a **round point**.
- Problem:** Classify the **ancient compact** embedded solutions to the Curve shortening flow.


Ancient Convex solutions to the CSF

- The **curvature** κ of Γ_t evolves, in terms of its **arc-length** s , by

$$\kappa_t = \kappa_{ss} + \kappa^3.$$

- Examples of solutions in closed form:

- Type I**: the contracting circles.
- Type II**: the **Angenent ovals (paper clips)**. They are **not solitons**. As $t \rightarrow -\infty$ they may be visualized as two **grim reaper** solutions glued together.



The Classification of Ancient Convex solutions to the CSF

- **Theorem** (D., Hamilton, Sesum - 2010)
The **only** ancient convex solutions to the CSF are the contracting spheres or the Angenent ovals.
- **Proof:** It is based on:
 - i. two **monotonicity formulas** and
 - ii. the fact that at its **singular time** any solution becomes **circular** with **very sharp rates of convergence**.

Non-Convex ancient solutions

- **Question:** Do they exist **non convex compact** embedded solutions to the curve shortening flow ?
- **Angenent (2011):** Presents a **YouTube video** of an ancient solution to the CSF built out from one Yin-Yang spiral and one Grim Reaper.




- **S. Angenent:** is currently working on a rigorous construction of these solutions.

The Mean curvature flow

- Let M_t , $t \in (-\infty, T)$ be a smooth **ancient** compact solution of the **Mean curvature flow**

$$(MCF) \quad \frac{\partial F}{\partial t} = -H \nu$$




$H(p, t)$ is the **Mean curvature** and ν a choice of **unit normal**.

- Problem:** Understand **ancient compact** solutions M_t of the **Mean curvature flow**.

Ancient non-collapsed solutions to MCF

- Weimin Sheng and Xu-Jia Wang; Ben Andrews: Introduced an α -noncollapsed condition.


$$B = B_{\frac{\alpha}{H(p)}}$$



- Haslhofer & Kleiner (2013):
Ancient compact + α -noncollapsed MCF solution \Rightarrow convex.
- convex compact + self-similar MCF solution \Rightarrow sphere.
- Ancient ovals: Any compact and α -noncollapsed solution to MCF which is not self-similar.
- Other ancient solutions to MCF: compact and collapsed.

Ancient MCF ovals


- **Problem:** Provide the **classification** of all **Ancient ovals**.
- **B. White (2003):** Existence of certain **Ancient ovals** with $O(k) \times O(l)$ symmetry. We call them **White ancient ovals**.
- **Haslhofer & Hershkovits (2013):** Give more details in the existence proof of the White Ancient ovals.
- **Angenent (2012):** establishes the **formal matched asymptotics** of all **Ancient ovals** as $t \rightarrow -\infty$.



- They are small perturbations of **ellipsoids**.

Unique asymptotics of Ancient MCF ovals

- S. Angenent, D., and N. Sesum (2015): All ancient ovals with $O(1) \times O(n)$ symmetry have **unique asymptotics** as $t \rightarrow -\infty$, and satisfy Angenent's **precise matched asymptotics**:



- **Geometric properties** $t \rightarrow -\infty$: **type II** ancient solutions

$$\text{diam}(t) \approx \sqrt{8|t| \log |t|} \quad \text{and} \quad H_{\max}(t) \approx \frac{\sqrt{\log |t|}}{\sqrt{2|t|}}.$$

- **The proof involves**: the analysis of the **linearized operator** at the cylinder, **Huisken's monotonicity formula** and carefully constructed **barriers** at the intermediate region.

Uniqueness of Ancient MCF ovals

- **Work in progress:** to establish such asymptotics in the **non-symmetric** case.
- **Next Step:** Establish the **uniqueness** of the **Ancient ovals**.
- **Conjecture 1:** The **Ancient ovals** with $O(l) \times O(k)$ symmetry are **uniquely** determined by their **asymptotics** at $t \rightarrow -\infty$.
- **Hence:** they are **unique** (up to dilation and translation in rescaled time).
- **Conjecture 2:** All **Ancient ovals** are $O(l) \times O(k)$ symmetric.

Eternal convex solutions

- Work in progress (with N. Sesum) Classify all eternal convex and complete solutions to the MCF.
- **Conjecture:** Eternal convex and entire graph solutions are translating solitons.
- **Haslhofer:** A uniformly two convex ($\lambda_1 + \lambda_2 \geq \beta H$) and α -noncollapsed translating soliton is necessarily a bowl soliton.
- Work in progress (with Davila, delPino, Wei): We construct non-convex entire graph translating solitons in dimensions $n \geq 8$.

Ancient compact solutions to the 2-dim Ricci flow

- Consider an ancient solution of the Ricci flow

$$(RF) \quad \frac{\partial g_{ij}}{\partial t} = -2 R_{ij}$$

on a compact manifold M^2 which exists for all time $-\infty < t < T$ and becomes singular at time T .

- In dim 2, we have $R_{ij} = \frac{1}{2} R g_{ij}$, where R is the scalar curvature.
- Hamilton (1988), Chow (1991): After re-normalization, the metric becomes spherical at $t = T$.
- Problem:** Provide the classification of all ancient compact solutions.

Ancient compact solutions to the 2-dim Ricci flow

- Choose a parametrization $g_{S^2} = d\psi^2 + \cos^2 \psi d\theta^2$ of the limiting spherical metric.
- We parametrize our solution as $g(\cdot, t) = u(\cdot, t) g_{S^2}$.
- Then the (RF) becomes equivalent to the **fast-diffusion** equation:

$$u_t = \Delta_{S^2} \log u - 2, \quad \text{on } S^2 \times (-\infty, T).$$

- Provide the **classification** of all ancient solutions.

Examples of compact solutions on S^2


- **Type I** solution: the contracting spheres.



- **Type II** solution: the **King-Rosenau** solution of the form:

$$u(\psi, t) = [a(t) + b(t) \sin^2 \psi]^{-1}, \quad t < T.$$

As $t \rightarrow -\infty$ the King-Rosenau solution looks like two **cigar solitons** glued together.



The classification result

Theorem: (D., Hamilton, Sesum - 2012)

The only **ancient** solutions to the Ricci flow on S^2 are the **contracting spheres** and the **King-Rosenau** solutions.

Proof: combines geometric arguments and PDE techniques.

- i. a **monotonicity formula** and uniform a priori $C^{1,\alpha}$ estimates that allow us to pass to the limit as $t \rightarrow -\infty$.
- ii. **geometric arguments** that allow us to **classify** the **backward limit** as $t \rightarrow -\infty$.
- iii. **maximum principle** arguments that allow us to **characterize** the **King-Rosenau** solutions among type II solutions.
- iv. an **isoperimetric inequality** that allows us to **characterize** the **contracting spheres** among type I solutions.

The 3 dimensional Ricci flow - Open problems

- **3-dim Ricci flow:** The analogue of the 2-dim King-Rosenau solutions have been shown to exist by **G. Perelman**. They are not given in closed form, they are **type II** and **k-noncollapsed**.
- Other **collapsed** compact solutions in closed form have been found by **V.A. Fateev** in a paper dated back to 1996.
- **Conjecture:** The only **k-noncollapsed** ancient and compact solutions to the 3-dim Ricci flow are the contracting spheres and the **Perelman** solutions.

Ancient solutions to the Yamabe flow

- We will conclude by discussing **ancient solutions** $g = g_{ij}$ of the **Yamabe flow** on S^n , $n \geq 3$.
- The Yamabe flow may be viewed as the higher dimensional analogue of the 2-dim Ricci flow.
- It is the evolution of metric $g(\cdot, t)$ **conformally equivalent** to the standard metric on S^n by

$$\frac{\partial g}{\partial t} = -Rg \quad \text{on } -\infty < t < T$$

where R denotes the **scalar curvature** of g .

- **Question:** Is it possible to provide the **classification** of all such ancient solutions ?

The Yamabe flow - Background

- Let (M^n, g_0) , $n \geq 3$ be a compact manifold without boundary. The scalar curvature R of a metric $g = v^{\frac{4}{n-2}} g_0$ conformal to g_0 is given by

$$R = -v^{-\frac{n+2}{n-2}} (c_n \Delta_{g_0} v - R_0 v)$$

where R_0 denotes the scalar curvature of g_0 .

- R. Hamilton (1989): introduced the Yamabe flow as a parabolic approach to resolve the Yamabe problem.
- S. Brendle (2007): convergence of the normalized flow to a metric of constant scalar curvature (up to a mild technical assumption for $\dim n \geq 6$).
- Previous important works: Hamilton '89, Chow '92, Ye '94, Schwetlick-Struwe '2003.

Ancient solutions to the Yamabe flow on S^n

- Let $g = v^{\frac{4}{n-2}} g_{S^n}$ be an **ancient** solution to the **Yamabe flow**, which is conformal to the standard metric on S^n .
- The function v evolves by the **fast diffusion** equation

$$\left(v^{\frac{n+2}{n-2}}\right)_t = \Delta_{S^n} v - c_n v \quad \text{on } S^n \times (-\infty, T).$$

- Let $g = \bar{v}^{\frac{4}{n-2}} g_{\mathbb{R}^n}$ after stereographic projection. Then,

$$\left(\bar{v}^{\frac{n+2}{n-2}}\right)_t = \Delta \bar{v} \quad \text{on } \mathbb{R}^n \times (-\infty, T).$$


The King Solutions

- J.King (1993): discovered **non-self similar type I** ancient compact solutions to the (YF) on S^n in closed form.
- **King solutions:** $g = \hat{v}_K(\cdot, t)^{\frac{4}{n-2}} g_{\mathbb{R}^n}$, where

$$\hat{v}_K(x, t) = (a(t) + 2b(t)|x|^2 + a(t)|x|^4)^{-\frac{n-2}{4}}, \quad x \in \mathbb{R}^n.$$

- As $t \rightarrow -\infty$ they converge (after rescaling) to two **Barenblatt** type self-similar solutions (**shrinking solitons**) joined by a long **cylindrical neck**.

$t \approx -\infty$




Ancient solutions to the Yamabe flow on S^n

- Question 1:
Are the contracting spheres and the King solutions the only examples of type I ancient solutions ?
- Question 2:
Are there any type II ancient solutions ?

New Type I solutions to the Yamabe flow

- **Recent work:** (D., del Pino, J. King and N. Sesum - 2015)
There exist **infinite many** other **type I** ancient solutions.
- As $t \rightarrow -\infty$ they look as two **self-similar solutions** v_λ, v_μ connected by a **cylinder** and moving with **speeds** $\lambda > 0, \mu > 0$.



- Our solutions are **not given in closed** form but we show **sharp asymptotics**.
- In **similar spirit** to the work by Hamel and Nadirashvili (1999) where they construct a five parameter family of **ancient** solutions for the **KPP equation**

$$u_t = u_{xx} + f(u), \quad x \in R.$$

Shrinking solitons with cylindrical behavior

- We look for rotationally symmetric **shrinking solitons** of the (YF) expressed in **cylindrical coordinates** $g = v^{\frac{4}{n-2}} g_{cyl}$.
- $v(x, \tau)$ satisfies (after a type I rescaling) the equation:

$$(*) \quad \left(v^{\frac{n+2}{n-2}}\right)_{\tau} = v_{xx} - v + v^{\frac{n+2}{n-2}}.$$


- **Shrinking solitons (or traveling waves):** $\forall \lambda \geq 1$ there exist a solution $v_{\lambda} = V_{\lambda}(x - \lambda\tau)$ of $(*)$ with **cylindrical** behavior

$$V_{\lambda}(x) \approx 1 - C_{\lambda} e^{-\gamma_{\lambda} x}, \quad \text{as } x \rightarrow +\infty.$$


- **Theorem:** (D., J. King and N. Sesum)
 L^1 stability of the traveling wave solutions v_{λ} .

Shrinking solitons with cylindrical behavior


- Consider **shrinking solitons** in cylindrical coordinates and after a type I scaling.
- Traveling wave to the right:** $v_{\lambda,h} = V_{\lambda}(x - \lambda \tau + h)$



- Traveling wave to the left:** $\bar{v}_{\mu,h'} = V_{\mu}(-x - \mu \tau + h')$




- Cylinder:** $\xi_k(\tau) \approx 1 - k e^{\tau/2}$, as $\tau \rightarrow -\infty$.



New type I ancient solutions

- **Theorem:** (D., del Pino, J. King and N. Sesum - 2016)

There exist a five parameter family $v_{\lambda,\mu,h,h',k}$ of **type I** ancient solutions of the **Yamabe flow** on $S^n \times (-\infty, T)$.



- In terms of the **pressure function** $f := v^q$, $q := -\frac{4}{n-2}$ it satisfies:

$$v_{\lambda,\mu,h,h',k}^q \approx v_{\lambda,h}^q(x, \tau) + \xi_k(\tau)^q + \bar{v}_{\mu,h'}^q - 2.$$

New type I ancient solutions - Sketch of proof

- **Hamel and Nadirashvili (1999)**: Construct a five parameter family of **ancient** solutions for $u_t = u_{xx} + f(u)$ by estimating the error of approximation using **heat kernel** representation.
- We construct precise barriers for the **pressure function**

$$f := v_{\lambda,\mu,h,h',k}^q, \quad q := -\frac{4}{n-2} :$$

$$w_{\lambda,\mu,h,h',k}^- := \max \left(v_{\lambda}^q(x - \lambda\tau + h), \xi_k^q(\tau), v_{\mu}^q(-x - \mu\tau + h') \right)$$

$$\begin{aligned} w_{\lambda,\mu,h,h',k}^+ &:= v_{\lambda}^q(x - \lambda\tau (1 - c e^{\frac{p-1}{p}\tau}) + h) + \xi_k^q(\tau) \\ &\quad + v_{\mu}^q(-x - \mu\tau (1 - c e^{\frac{p-1}{p}\tau}) + h') - 2 \end{aligned}$$

where $c = c(p)$.

- We have

$$w_{\lambda,\mu,h,h',k}^- \leq v_{\lambda,\mu,h,h',k}^q \leq w_{\lambda,\mu,h,h',k}^+, \quad -\infty < \tau < +\infty.$$

Ancient towers of moving bubbles - type II solutions

- **Question:** Are there any **type II** ancient solutions to (YF) ?
- **D., del Pino and Sesum (2013):**
We construct a class of **ancient solutions** of the **Yamabe flow** on S^n which (after re-normalization) converge as $t \rightarrow -\infty$ to a **tower of n-spheres**. They are rotationally symmetric.



- The **curvature operator** in these solutions **changes sign** and they are of **type II**.
- Our construction also holds for **any number of bubbles**.



Discussion on parabolic gluing methods

- Our construction may be viewed as a **parabolic analogue** of the **elliptic gluing** technique.
- **Elliptic gluing**: pioneering works by **Kapouleas '90 -'95** and works **Mazzeo, Pacard, Pollack, Ulhenbeck**.
- **Brendle & Kapouleas (2014)**: construct new **ancient compact** solutions to the **4-dim Ricci flow** by parabolic gluing.
- **Future research direction**: apply parabolic gluing on other geometric flows.

Conclusion

- We discussed **ancient solutions** to **geometric parabolic PDE**.
- Typical examples are either **solitons** or other **special solutions** obtained from the **gluing** as $t \rightarrow -\infty$ of solitons.
- The only existing classification results heavily rely on knowing the **exact form** of these ancient solutions.
- **Future research direction:** develop new techniques that allow us to **characterize** and **construct** other types of ancient or eternal solutions.