

# Strong arithmetic equivalence

Andrew V. Sutherland

Massachusetts Institute of Technology

June 23, 2017

Arithmetic Geometry, Cryptography, and Coding Theory

Centre International de Recontres Mathématiques

# Arithmetic equivalence

## Definition

Number fields  $K_1$  and  $K_2$  are **arithmetically equivalent** if  $\zeta_{K_1}(s) = \zeta_{K_2}(s)$ . The fields  $K_1 \sim K_2$  must have the same degree and Galois closure  $L$ .

Let  $G := \text{Gal}(L/\mathbb{Q})$ ,  $H_1 := \text{Gal}(L/K_1)$ , and  $H_2 := \text{Gal}(L/K_2)$ .

## Definition

A **Gassmann triple**  $(G, H_1, H_2)$  consists of finite groups  $H_1, H_2 \leq G$  that satisfy  $\#(H_1 \cap C) = \#(H_2 \cap C)$  for every  $G$ -conjugacy class  $C$ .

## Theorem (Gassmann 1926)

$K_1 \sim K_2$  if and only if  $(G, H_1, H_2)$  is a Gassmann triple.

$K_1 \simeq K_2$  if and only if  $H_1$  and  $H_2$  are conjugate in  $G$ .

## Examples

Let  $G = \mathrm{GL}_2(\mathbb{F}_3)$ , let  $H_1 = \{ \begin{pmatrix} * & * \\ 0 & 1 \end{pmatrix} \in G \}$ , and let  $H_2 = \{ \begin{pmatrix} 1 & * \\ 0 & * \end{pmatrix} \in G \}$ . Then  $(G, H_1, H_2)$  is a non-trivial Gassmann triple (de Smit, ANTS III).

Let  $E/\mathbb{Q}$  be an elliptic curve with surjective mod-3 Galois image and let  $L = \mathbb{Q}(E[3])$ . then  $\mathrm{Gal}(L/\mathbb{Q}) \simeq G$  and the fields  $K_1 := L^{H_1}$  and  $K_2 = L^{H_2}$  are non-conjugate arithmetically equivalent number fields of degree 8.

This example generalizes: one can replace  $p$  with any odd degree, and the matrix entry 1 can be replaced squares in  $\mathbb{F}_p$ ; the degree is  $2p + 2$ .

One can achieve degree 7 using similar subgroups of  $\mathrm{SL}_3(\mathbb{F}_2)$ , which is best possible (Bosma–de Smit, ANTS V, de Smit–Lenstra 2000).

The subgroups  $H_1$  and  $H_2$  need not be isomorphic; for example, take  $G \simeq [384, 3755]$ ,  $H_1 \simeq [16, 3]$ ,  $H_2 \simeq [16, 10]$  (in GAP notation).

## Gassmann triples in other contexts

Gassmann triples  $(G, H_1, H_2)$  arise in many other contexts involving potentially non-isomorphic objects with the same “zeta function”:

- If  $\pi: M \rightarrow M_0$  is a normal finite Riemannian covering with deck group  $G$  then  $M/H_1$ , and  $M/H_2$  are isospectral (Sunada 1985).
- If  $\Gamma$  is a finite graph with  $G = \text{Aut}(\Gamma)$  then  $\Gamma/H_1$  and  $\Gamma/H_2$  are isospectral (Halbeisen–Hungerbühler 1995).
- If  $X/k$  is a projective curve with  $G = \text{Aut}(X)$ , then  $X/H_1$  and  $X/H_2$  have isogenous Jacobians (Prasad–Rajan 2003).
- If  $\pi: X \rightarrow Y$  is a Galois étale cover of  $k$ -varieties then  $X/H_1$  and  $X/H_2$  have isomorphic Chow motives (Arapura et al. 2017).

Unlike the number field case, non-trivial Gassmann triples may give rise to isomorphic objects (imposing further conditions prevents this).

# Characterizations of Gassmann triples

Let  $[G/H]$  be the transitive  $G$ -set consisting of cosets of  $H$ .  
Let  $\chi_H: G \rightarrow \mathbb{Z}$  be the permutation character  $g \mapsto \#[G/H]^g$ , and  
for  $K \leq G$  define  $\chi_H(K) := \#[G/H]^K$ . We then have

$$\chi_H(K) \neq 0 \iff K \leq_G H$$

(indeed,  $HgK = Hg \iff gKg^{-1} \subseteq H$ ).

## Proposition

For  $H_1, H_2 \leq G$  the following are equivalent:

- 1  $\#(H_1 \cap C) = \#(H_2 \cap C)$  for all  $C \in \text{conj}(G)$ .
- 2 There is a  $G$ -conjugacy preserving bijection  $H_1 \leftrightarrow H_2$ .
- 3  $\chi_{H_1}(K) = \chi_{H_2}(K)$  for all cyclic subgroups  $K \leq G$  (or all  $K \leq H_1, H_2$ ).
- 4  $\mathbb{Q}[G/H_1] \simeq \mathbb{Q}[G/H_2]$  (as  $\mathbb{Q}[G]$  modules).

# How strong is arithmetic equivalence?

Let  $K_1$  and  $K_2$  be arithmetically equivalent number fields.

## Theorem (Perlis 1977)

*The number fields  $K_1$  and  $K_2$  have the same degree, discriminant, signature, and roots of unity.*

The analytic class number formula

$$\lim_{s \rightarrow 1^+} (s-1)\zeta_K(s) = \frac{2^{r_1} (2\pi)^{r_2} h_K R_K}{w_K |D_K|^{1/2}}$$

implies  $R_{K_1} h_{K_1} = R_{K_2} h_{K_2}$  but the class numbers and regulators may differ.

There is a bijection of the places of  $K_1$  and  $K_2$  that preserves residue field degrees, but not necessarily ramification indices.

The adèle rings and idèle groups of  $K_1$  and  $K_2$  need not be isomorphic.

# Stronger notions of arithmetic equivalence

## Definition

Two number fields are **locally isomorphic** if there is a bijection of their places such that corresponding completions are isomorphic.

Locally isomorphic fields are arithmetically equivalent (Klingen 1998).

## Proposition (Iwasawa 1953)

*Two number fields  $K_1, K_2$  are locally isomorphic if and only if they have isomorphic rings of adèles  $\mathbb{A}_{K_1} \simeq \mathbb{A}_{K_2}$ .*

## Proposition (Linowitz–McReynolds–Miller 2017)

*Locally isomorphic number fields have isomorphic Brauer groups.*

But locally isomorphic fields may have distinct class numbers, as happens with  $\mathbb{Q}(\sqrt[8]{-33})$  and  $\mathbb{Q}(\sqrt[8]{-33 \cdot 16})$  (de Smit–Perlis, 1994).

## Local integral equivalence

A finite group  $K$  is  *$p$ -cyclic* (or  *$p$ -hypoelementary*) if the quotient of  $K$  by the intersection of its  $p$ -Sylow subgroups (its  $p$ -core) is cyclic.

### Proposition

Let  $p$  be a prime. For  $H_1, H_2 \leq G$  the following are equivalent:

- $\chi_{H_1}(K) = \chi_{H_2}(K)$  for all  $p$ -cyclic  $K \leq G$  (or all  $K \leq H_1, H_2$ );
- $\mathbb{Z}_p[G/H_1] \simeq \mathbb{Z}_p[G/H_2]$ ;
- $\mathbb{F}_p[G/H_1] \simeq \mathbb{F}_p[G/H_2]$ ;
- $\det(\text{Hom}_{\mathbb{Z}[G]}(\mathbb{Z}[G/H_1], \mathbb{Z}[G/H_2])) \notin p\mathbb{Z}$ .

If  $\mathbb{Z}_p[G/H_1] \simeq \mathbb{Z}_p[G/H_2]$  for all  $p$  we have *local integral equivalence*.

### Theorem (Perlis 1978)

Number fields  $K_1, K_2$  corresponding to a locally integrally equivalent  $H_1, H_2 \leq G$  have isomorphic class groups.



# Integral equivalence

## Definition

Subgroups  $H_1, H_2 \leq G$  are **integrally equivalent** if  $\mathbb{Z}[G/H_1] \simeq \mathbb{Z}[G/H_2]$ .

Let  $H_1, H_2 \leq G$  have index  $n$ , let  $\rho_1, \rho_2: G \rightarrow S_n$  be the representations corresponding to the permutation modules  $\mathbb{Z}[G/H_1], \mathbb{Z}[G/H_2]$ .

Fix an ordering of  $[G/H_1]$  and  $[G/H_2]$ . We may represent elements of  $\text{Hom}_{\mathbb{Z}[G]}(\mathbb{Z}[G/H_1], \mathbb{Z}[G/H_2]) \not\subseteq p\mathbb{Z}$  by matrices  $M \in \mathbb{Z}^{n \times n}$  that satisfy

$$M_{ij} = M_{\rho_1(g)(i), \rho_2(g)(j)} \quad \text{for all } g \in G.$$

Our two notions of integral equivalence are distinguished by:

- **local integral equivalence**:  $\gcd(\det(M_1), \dots, \det(M_r)) = 1$  for some  $M_1, \dots, M_r \in \text{Hom}_{\mathbb{Z}[G]}(\mathbb{Z}[G/H_1], \mathbb{Z}[G/H_2])$ .
- **global integral equivalence**:  $\det(M) = \pm 1$  for some  $M \in \text{Hom}_{\mathbb{Z}[G]}(\mathbb{Z}[G/H_1], \mathbb{Z}[G/H_2])$ .

# What we know about integral equivalence

## Theorem (Prasad 2017)

Let  $\pi: X \rightarrow Y$  be a Galois cover of nice curves over  $k$  with Galois group  $G$ . If  $H_1, H_2 \leq G$  are integrally equivalent then  $\text{Jac}(X/H_1) \simeq \text{Jac}(X/H_2)$ .

**Remark:** Infinite families of non-isomorphic curves of low genus with isomorphic Jacobians were previously known (Howe 2005).

Essentially only one non-trivial example of integral equivalence is known:  $G = \text{PSL}_2(\mathbb{F}_{29})$  with  $H_1, H_2 \simeq A_5$  non-conjugate of index 203.

This example is due to Leonard Scott, who proved it by explicitly exhibiting  $M \in \text{Hom}_{\mathbb{Z}[G]}(\mathbb{Z}[G/H_1], \mathbb{Z}[G/H_2]) \subseteq \mathbb{Z}^{203 \times 203}$  with  $\det M = 1$ .

Similar triples exist for  $p \equiv \pm 29 \pmod{120} \dots$

$\dots$  but for  $p = 149$  we already have to work with  $M \in \mathbb{Z}^{27565 \times 27565}$ .

# What we don't know about integral equivalence

**Question 1:** Must integrally equivalent  $H_1, H_2 \leq G$  be isomorphic?  
How about locally integrally equivalent  $H_1, H_2 \leq G$ ?

Both necessarily hold if  $G = \mathrm{PSL}_2(\mathbb{F}_p)$ . In fact, Gassmann equivalent subgroups of  $\mathrm{PSL}_2(\mathbb{F}_p)$ ,  $\mathrm{SL}_2(\mathbb{F}_p)$ ,  $\mathrm{GL}_2(\mathbb{F}_p)$  are isomorphic (S 2016).

Scott's triple gives rise to infinitely many arithmetically equivalent number non-isomorphic number fields of degree 203 that are also locally isomorphic, hence have isomorphic adèle rings.

But (as noted by Prasad), it is not clear that integral equivalence alone guarantees that we will get locally isomorphic number fields.

**Question 2:** Do locally integrally equivalent  $H_1, H_2 \leq G$  give rise to locally isomorphic number fields? If not, does integral equivalence?

# Solvable integral equivalence

## Definition

Subgroups  $H_1, H_2 \leq G$  are **solvably equivalent** if  $\chi_{H_1}(K) = \chi_{H_2}(K)$  for all solvable subgroups  $K \leq G$ .

Like integral equivalence, solvable equivalence obviously implies local integral equivalence (hence isomorphic class groups).

## Proposition

*Number fields  $K_1, K_2$  corresponding to solvably equivalent  $H_1, H_2 \leq G$  are arithmetically equivalent, locally isomorphic, and have the same class number. Moreover, there is a bijection of the places of  $K_1$  and  $K_2$  that preserves residue degrees and ramification indices.*

**Remark:** Solvable equivalence is stronger than necessary

# Results

## Proposition

*There are infinitely many non-isomorphic pairs of degree 32 number fields arising from locally (not globally) integrally equivalent  $H_1, H_2 \leq G$ .*

## Proposition

*There are infinitely many non-isomorphic pairs of degree 96 number fields arising from solvably (not integrally) equivalent  $H_1, H_2 \leq G$ .*

These results are effective; we can construct explicit examples.

The fact that  $\mathrm{PSL}_2(\mathbb{F}_p)$  contains non-conjugate solvably equivalent subgroups  $H_1, H_2$  for all primes  $p \equiv \pm 29 \pmod{120}$  implies that there are infinitely many non-isomorphic pairs of number fields arising from infinitely many solvably equivalent  $H_1, H_2 \leq G$ , but the degrees of these fields is at least 203 and they hard to construct explicitly.

## First example

An exhaustive search of the 11,759,892 groups of order less than 1024 finds exactly 74 groups  $G$  that contain non-conjugate locally integrally equivalent subgroups  $H_1, H_2$ .

The smallest two have GAP ids [384, 18050] and [384, 18046], isomorphic to transitive permutation groups  $32T9403$  and  $32T9408$ . Both are 2-extensions of  $D_4 \times S_4$ , which makes it easy to construct explicit examples.

For instance, the polynomials

$$\begin{aligned}x^{32} + 12x^{28} + 72x^{24} + 120x^{20} - 234x^{16} + 108x^{12} + 396x^8 - 432x^4 + 81, \\x^{32} - 12x^{28} + 72x^{24} - 120x^{20} - 234x^{16} - 108x^{12} + 396x^8 + 432x^4 + 81\end{aligned}$$

both have Galois group  $G = 32T9403$ . They define non-isomorphic fields  $K_1, K_2$  corresponding to locally integrally equivalent  $H_1, H_2 \leq G$ .

## First example (continued)

We can view each  $M \in \text{Hom}_{\mathbb{Z}[G]}(\mathbb{Z}[G/H_1], \mathbb{Z}[G/H_2])$  as  $32 \times 32$  matrix with entries  $x_1, \dots, x_8 \in \mathbb{Z}$ . corresponding to the decomposition of  $G$  into eight double cosets  $H_1 g H_2$ . A (non-trivial) calculation finds that

$$\begin{aligned} \det M = & - (2(x_2 - x_3)^2 + 3(x_5 - x_6)^2)^8 \\ & \cdot (2(x_1 - x_4) + (x_5 + x_6 - 2x_7))^6 \\ & \cdot (2(x_1 + x_2 + x_3 + x_4) - (x_5 + x_6 + 2x_7 + 4x_8))^3 \\ & \cdot (2(x_1 - x_2 - x_3 + x_4) - (x_5 + x_6 + 2x_7 - 4x_8))^3 \\ & \cdot (2(x_1 - x_4) - 3(x_5 + x_6 - 2x_7))^2 \\ & \cdot (2(x_1 + x_2 + x_3 + x_4) + 3(x_5 + x_6 + 2x_7 + 4x_8)) \\ & \cdot (2(x_1 - x_2 - x_3 + x_4) + 3(x_5 + x_6 + 2x_7 - 4x_8)). \end{aligned}$$

One can choose the  $x_i$  so that  $\det M = 2^{32}$ , and so that  $\det M = 3^{12}$ . Thus  $H_1$  and  $H_2$  are locally integrally equivalent, but they not integrally equivalent because there is no choice of the  $x_i$  for which  $\det M = \pm 1$ .

This negatively answers a question of Guralnick–Weiss from 1993.

## Second example

Let  $G = 16T1654$  of order 5760. It contains non-conjugate  $H_1, H_2 \simeq A_5$  of index 96 such that every proper subgroup of  $H_1$  is a proper subgroup of  $H_2$ .

The group  $G$  is the Galois group of an extension of  $\mathbb{Q}[T]$ ; Hilbert irreducibility gives infinitely many examples of corresponding number fields, including:

$$x^{16} - 2x^{15} + 3x^{14} - 16x^{13} + 18x^{12} - 10x^{10} + 40x^9 - 39x^8 + 54x^7 + 23x^6 + 16x^5 - 140x^4 - 188x^3 - 28x^2 + 104x - 4,$$

Each  $M \in \text{Hom}_{\mathbb{Z}[G]}(\mathbb{Z}[G/H_1], \mathbb{Z}[G/H_2])$  has entries  $x_1, x_2, x_3, x_4, x_5 \in \mathbb{Z}$ , and

$$\begin{aligned} \det M = & - (5x_1 + 6x_2 + 10x_3 + 15x_4 + 60x_5) \\ & \cdot (x_1 - 6x_2 - 10x_3 + 3x_4 + 12x_5)^5 \\ & \cdot (3x_1 + 2x_2 - 2x_3 - 7x_4 + 4x_5)^{15} \\ & \cdot (3x_1 - 2x_2 + 2x_3 + x_4 - 4x_5)^{30} \\ & \cdot (x_1 + 2x_2 - 2x_3 + 3x_4 - 4x_5)^{45} \end{aligned}$$

No assignment of  $x_1, x_2, x_3, x_4, x_5 \in \mathbb{Z}$  makes every factor in  $\det M$  equal to  $\pm 1$ , so  $H_1$  and  $H_2$  are not integrally equivalent.