

FROM COMBINATORIAL GAMES TO SHAPE-SYMMETRIC MORPHISMS

Michel Rigo


<http://www.discmath.ulg.ac.be/>

<http://orbi.ulg.ac.be/>

Jean-Morlet chair: Shigeki Akiyama

Research school: Tiling Dynamical System, 20–24 November 2017

Université
de Liège



TENTATIVE OUTLINE OF THE TALKS

Lecture 1:

- ▶ Starting from CGT
- ▶ Recap on k -automatic sequences

Lecture 2:

- ▶ Moving to multidimensional sequences
- ▶ Motivations from CGT: Wythoff's \mathcal{P} -positions
- ▶ Maes' shape-symmetric morphisms

Lecture 3:

- ▶ Link with (abstract) numeration systems
- ▶ Back to CGT and Presburger definable sets, work in progress

NOTATION AND CONVENTIONS

- ▶ MSDF most significant digit first (on the left)
- ▶ $k \geq 2$ is an integer
- ▶ $\text{rep}_2(81) = 1010001$; rep_k canonical base- k *representation*
- ▶ numerical *value*

$$\text{val}_k(d_\ell \cdots d_0) = \sum_{j=0}^{\ell} d_j k^j$$

- ▶ extension to another system S : rep_S and val_S
- ▶ $\mathbb{N} = \{0, 1, 2, \dots\}$
- ▶ an infinite word $w : \mathbb{N} \rightarrow A$ (indexing starts with 0)

LET'S TRY TO MIX TWO TOPICS

↪ Motivations from *Combinatorial Game Theory* (CGT)

- ▶ independent community of researchers
- ▶ use tools from number theory, numeration systems, continued fractions, combinatorics on words, cellular automata, etc.

Some general references in CGT:

- ▶ A. Siegel, *Combinatorial game theory*, Graduate Studies in Mathematics, **146**, AMS (2013). [1]
- ▶ E. R. Berlekamp, J. H. Conway, R. K. Guy, *Winning ways for your mathematical plays*, A K Peters, Ltd., (2001). [2]
- ▶ T. S. Ferguson, *Game Theory*, UCLA. [3]

Wythoff's game [5] or, *the Queen* ♔ goes to $(0, 0)$

- ▶ **two players** playing alternatively;
- ▶ the player *unable to move* loses the game (**Normal play**);
- ▶ two piles of token;
- ▶ Nim rule [6]: remove a positive number of token from one pile
♙

$$\text{Moves} = \{(i, 0), (0, i) \mid i \geq 1\}.$$

- ▶ Wythoff's rule: remove *simultaneously* the same number of token from both piles


$$\text{Moves} = \{(i, 0), (0, i), (i, i) \mid i \geq 1\}.$$

♔ (6, 3)

6, 3


0, 0

♔ (6, 3)




$$\text{crown} (6, 3) \xrightarrow{A} (4, 1)$$

6, 3



$$\text{♔ } (6, 3) \xrightarrow{A} (4, 1) \xrightarrow{B} (2, 1)$$



$$\text{♔ } (6, 3) \xrightarrow{A} (4, 1) \xrightarrow{B} (2, 1) \xrightarrow{A} (1, 0)$$

(6, 3)

(2, 1)

(4, 1)

(0, 0) — (1, 0)

$$\text{♔ } (6, 3) \xrightarrow{A} (4, 1) \xrightarrow{B} (2, 1) \xrightarrow{A} (1, 0) \xrightarrow{B} (0, 0)$$

(6, 3)

(2, 1)

(4, 1)

(0, 0)

(1, 0)

WINNING AND LOSING POSITIONS

STATUS \mathcal{N} (NEXT) OR \mathcal{P} (PREVIOUS)


A position is \mathcal{P} , if all its options are \mathcal{N} ;

A position is \mathcal{N} , if there exists an option in \mathcal{P} .

If the *game-graph*:

- ▶ vertices = positions
- ▶ edges = available options


is acyclic, then every position is either \mathcal{N} , or \mathcal{P} .




REMARK (GRAPH-THEORETIC NOTION)

- ▶ The set of \mathcal{P} -positions is the *kernel* of the game-graph [4].
- ▶ The game-graph grows exponentially.


A *winning strategy* is a map from \mathcal{N} to \mathcal{P} assigning to every winning position in \mathcal{N} an available option in \mathcal{P} .




\mathcal{P} -positions and \mathcal{N} -positions for Wythoff's game.




\mathcal{P} -positions and \mathcal{N} -positions for Wythoff's game.




\mathcal{P} -positions and \mathcal{N} -positions for Wythoff's game.



\mathcal{P} -positions and \mathcal{N} -positions for Wythoff's game.



\mathcal{P} -positions and \mathcal{N} -positions for Wythoff's game.



\mathcal{P} -positions and \mathcal{N} -positions for Wythoff's game.

General questions

- ▶ Characterize the set of \mathcal{P} -positions?
- ▶ Is it computationally hard to determine these positions?
- ▶ Compute a winning strategy.

THEOREM (W. A. WYTHOFF, 1907 [5])

(x, y) is a \mathcal{P} -position if and only if it is of the form

$$(\lfloor n\varphi \rfloor, \lfloor n\varphi^2 \rfloor), \quad \text{for some } n$$

where φ is the Golden mean.

Several other characterizations are known. \rightsquigarrow Exercise: try to prove the above result (we will do it for Nim).

General questions

- ▶ Characterize the set of \mathcal{P} -positions?
- ▶ Is it computationally hard to determine these positions?
- ▶ Compute a winning strategy.

THEOREM (W. A. WYTHOFF, 1907 [5])

(x, y) is a \mathcal{P} -position if and only if it is of the form

$$(\lfloor n\varphi \rfloor, \lfloor n\varphi^2 \rfloor), \quad \text{for some } n$$

where φ is the Golden mean.

Several other characterizations are known. \rightsquigarrow Exercise: try to prove the above result (we will do it for Nim).

WINNING STRATEGY FOR NIM : XOR OR NIM-SUM

$$7 \oplus 2 \oplus 9 = 12$$

$$\begin{array}{rcccc} & & 1 & 1 & 1 \\ & & & 1 & 0 \\ 1 & 0 & 0 & 1 & \\ \hline 1 & 1 & 0 & 0 & \end{array}$$

THEOREM (L. BOUTON, 1902 [6])

*For the game of Nim with n piles of token,
a position (x_1, \dots, x_n) is in \mathcal{P} if and only if*

$$\bigoplus_{i=1}^n x_i = 0.$$

Exercise: (\mathbb{N}, \oplus) is an abelian group.

Proof.

1) $(0, \dots, 0)$ is in \mathcal{P} .

2) If a position has a zero Nim-sum,
all its options have a non-zero Nim-sum

$$\begin{array}{cccc} 1 & 0 & 1 & 1 \\ & & 1 & 0 \\ 1 & 0 & 0 & 1 \\ \hline 0 & 0 & 0 & 0 \end{array}$$

On a single row, one changes at least one digit \rightsquigarrow non-zero sum.

3) If a position has a non-zero Nim-sum,
there is an option having a zero Nim-sum

$$\begin{array}{rrrr} & 1 & 1 & 1 \\ & & 1 & 0 \\ 1 & 0 & 0 & 1 \\ \hline 1 & 1 & 0 & 0 \end{array}$$



3) If a position has a non-zero Nim-sum,
there is an option having a zero Nim-sum

$$\begin{array}{cccc} & 1 & 1 & 1 \\ & & 1 & 0 \\ \textcolor{red}{1} & \textcolor{red}{0} & \textcolor{red}{0} & \textcolor{red}{1} \\ \hline 1 & 1 & 0 & 0 \end{array}$$



3) If a position has a non-zero Nim-sum,
there is an option having a zero Nim-sum

$$\begin{array}{cccc} & 1 & 1 & 1 \\ & & 1 & 0 \\ 0 & 1 & 0 & 1 \\ \hline 0 & 0 & 0 & 0 \end{array}$$



GRUNDY FUNCTION (FOR SUMS OF GAMES)

DEFINITION

Let $S \subset \mathbb{N}$. **MeX** (minimum excluded value) of $S = \min \mathbb{N} \setminus S$.

Let G be a combinatorial game and x be a position.

The *Grundy function* is given by

$$\mathcal{G}(x) = \text{MeX}(\mathcal{G}(\text{Opt}(x))).$$

$$\text{MeX}\{0, 1, 3, 5\} = 2, \quad \text{MeX}\{2, 3, 6\} = 0.$$

We have a characterization of the **P**-positions:

IMPORTANT OBSERVATION

$\text{MeX } \emptyset = 0$, thus $\mathcal{G}(x) = 0$ iff x is in **P**.

GRUNDY FUNCTION (FOR SUMS OF GAMES)

THEOREM (SPRAGUE–GRUNDY [3])

Let G_i be combinatorial games with \mathcal{G}_i as Grundy function, $i = 1, \dots, n$. Then $G_1 + \dots + G_n$ has Grundy function

$$\mathcal{G}(x_1, \dots, x_n) = \mathcal{G}_1(x_1) \oplus \dots \oplus \mathcal{G}_n(x_n).$$

APPLICATION

Let's play on four boards simultaneously:

- ▶ G_1 Nim $\mathcal{G}_1(2, 5) = 7$
- ▶ G_2 Wythoff $\mathcal{G}_2(3, 4) = 2$
- ▶ G_3 Nim on three piles $\mathcal{G}_3(8, 7, 6) = 9$
- ▶ G_4 Wythoff $\mathcal{G}_4(3, 9) = 12$

Should you start? Just compute whether $7 \oplus 2 \oplus 9 \oplus 12$ is 0 or not?

General questions (one more)

- ▶ Characterize the set of \mathcal{P} -positions?
- ▶ Is it computationally hard to determine these positions?
- ▶ Compute a winning strategy.

Thanks to Sprague–Grundy theorem, we have an extra motivation:

- ▶ *Compute the Grundy function of all positions.*

For the game of Nim, first few values of $(x, y) \mapsto \mathcal{G}_N(x, y) = x \oplus y$

\vdots	\vdots										\ddots
9	9	8	11	10	13	12	15	14	1	0	
8	8	9	10	11	12	13	14	15	0	1	
7	7	6	5	4	3	2	1	0	15	14	
6	6	7	4	5	2	3	0	1	14	15	
5	5	4	7	6	1	0	3	2	13	12	
4	4	5	6	7	0	1	2	3	12	13	
3	3	2	1	0	7	6	5	4	11	10	
2	2	3	0	1	6	7	4	5	10	11	
1	1	0	3	2	5	4	7	6	9	8	
0	0	1	2	3	4	5	6	7	8	9	...
	0	1	2	3	4	5	6	7	8	9	...

\rightsquigarrow Exercises 21 and 22 in Section 16.6, p.451, Allouche–Shallit'03 [7].

What can be said about the structure of this table?

k -AUTOMATIC AND k -REGULAR SEQUENCES

There are various characterizations of k -automatic sequences:

- ▶ Image under a coding of a fixed point of a k -uniform morphism;
- ▶ Sequence of outputs of a DFAO fed with base- k expansions;
- ▶ Sets of indices defined by a first order formula in $\langle \mathbb{N}, +, V_k \rangle$;
- ▶ p -algebraic formal power series;
- ▶ Column of the space-time diagram of a cellular automaton;
- ▶ Finiteness of the k -kernel.


A. Cobham 1972 [8], Büchi–Bruyère theorem 1960–1985 [9], Christol–Kamae–Mendès France–Rauzy 1980 [12, 13], Rowland–Yassawi 2015 [14], S. Eilenberg 1974 [15, 11].

\rightsquigarrow see the survey by Bruyère, Hansel, Michaux and Villemaire [10].

Illustration 1/3

- Image under a coding of a fixed point of a k -uniform morphism;
- Sequence of outputs of a DFAO fed with base- k expansions;

$$f : \begin{cases} a \mapsto abc \\ b \mapsto cbc \\ c \mapsto bca \end{cases}$$



$$f^\omega(a) = abccbcbcabcacbcacbcacbcacbcbaabc \dots$$

$$g(f^\omega(a)) = 100000001001000001000001100 \dots$$

Let $f : A^* \rightarrow A^*$ such that $|f(a)| = k$ for all $a \in A$.




FIGURE: Constant length morphism $k = 4$.

$$\mathbf{x} = f^\omega(a) = x_0 x_1 x_2 \cdots x_q \cdots \cdots x_j \cdots \cdots$$

FUNDAMENTAL LEMMA (EXERCISE) [8]

Let j , $k^m \leq j < k^{m+1}$. We have $j = kq + r$, $k^{m-1} \leq q < k^m$ and $0 \leq r < k$. The symbol x_j is the $(r + 1)$ th symbol in $f(x_q)$.

$$k = 3$$
 a
$$a \quad b \quad c$$

$a \quad b \quad c \quad c \quad b \quad c \quad b \quad c \quad a$

$a \ b \ c \ c \ b \ c \ b \ c \ a \ b \ c \ a \ c \ b \ c \ b \ c \ a \ \dots$

0 1 2 3 4

↑
14

$14 = 3.4 + 2$ i.e. c is the third letter in $f(x_4) = f(b) = cbc$

$4 = 3.1 + 1$ i.e. x_4 is the second letter in $f(x_1) = f(b) = cbc$

$$\text{rep}_2(14) = 112.$$

$$k = 3$$
 a $a \quad b \quad c$

$a \quad b \quad c \quad c \quad b \quad c \quad b \quad c \quad a$

$a \ b \ c \ c \ b \ c \ b \ c \ a \ b \ c \ a \ c \ b \ c \ b \ c \ a \ \dots$

0 1 2 3 4

↑
14

$14 = 3.4 + 2$ i.e. c is the third letter in $f(x_4) = f(b) = cbc$


$4 = 3.1 + 1$ i.e. x_4 is the second letter in $f(x_1) = f(b) = cbc$

$$\text{rep}_2(14) = 112.$$

Exercise: prove that the *factor complexity* of a k -automatic word \mathbf{x} is in $\mathcal{O}(n)$. Hint: For $n \in (k^{\ell-1}, k^\ell]$, each factor of length n is inside $x(jk^\ell) \cdots x((j+2)k^\ell - 1)$ for some j . [8]

Illustration 2/3

- ▶ Sequence of outputs of a DFAO fed with base- k expansions;
- ▶ Sets of indices defined by a first order formula in $\langle \mathbb{N}, +, V_k \rangle$;



$$X = \{1, 2, 3, 4, 5, 6, 8, 9, 10, 12, 16, 17, 18, 20, 24, \dots\}$$

$$\text{rep}_2(X) = \{1, 10, 11, 100, 101, 110, 1000, 1001, 1010, 1100, \dots\}$$

$$X = \{n \in \mathbb{N} \mid \exists i, j \geq 0 : n = 2^i + 2^j\} \cup \{1\}$$

$$\psi(n) \equiv (n = 1) \vee (\exists i)(\exists j)(V_2(i) = i \wedge V_2(j) = j \wedge n = i + j).$$

Illustration 3/3

- Rowland–Yassawi: A sequence over a finite field \mathbb{F}_q of characteristic p is p -automatic if and only if it occurs as a column of the space-time diagram, with eventually periodic initial conditions, of a linear cellular automaton with memory over \mathbb{F}_q . [14]




FIGURE 1. Spacetime diagram of a linear cellular automaton with memory 12 containing the Thue–Morse sequence as a column.

DEFINITION [15]

The **k -kernel** of a sequence $\mathbf{x} = (x(n))_{n \geq 0}$ is a set of subsequences:

$$\text{Ker}_k(\mathbf{x}) = \{(x(k^i n + s))_{n \geq 0} \mid i \geq 0, 0 \leq s < k^i\}.$$

Alternative definition, introduce k operators of **k -decimation**, $r \in \{0, \dots, k-1\}$,

$$\partial_{k,r}((x(n))_{n \geq 0}) = (x(kn + r))_{n \geq 0}.$$

Thus the k -kernel is the set of sequences of the form






$$\partial_{k,r_1} \circ \dots \circ \partial_{k,r_m}((x(n))_{n \geq 0}).$$

REMARK

This corresponds to selecting a suffix in base- k expansions.

$$\text{val}_k(0^{i-m} r_m \dots r_1) = s.$$

Some of these subsequences; suffixes: $\varepsilon, 0, 1, 00, 01, 10, 11, 000$



S. EILENBERG, ALLOUCHE–SHALLIT THM. 6.6.2 [7]

A sequence is k -automatic IFF its k -kernel is finite.



\Rightarrow A DFAO is given. The transition monoid is finite $\leq \#Q^{\#Q}$

$$f_u : Q \rightarrow Q, q \mapsto q \cdot u$$

action of $\{0, \dots, k-1\}^*$ on the set of states Q ;



- ▶ reading $\varepsilon, 1, 2, 10, 11, 12, 100, 101, \dots$ provides us with the sequence $x(n)_{n \geq 0}$;
- ▶ reading $\varepsilon.u, 1.u, 2.u, 10.u, 11.u, 12.u, 100.u, 101.u, \dots$ provides us with the sequence of the k -kernel corresponding to $x(k^{|u|}n + \text{val}_k(u))_{n \geq 0}$.





		0	1	2	01	02	10	20	22	010	020	022	201	202	220	0220	2022
a	a	a	b	c	b	c	c	b	a	c	b	a	b	c	a	a	a
b	b	c	b	c	c	a	c	b	a	b	a	c	b	c	a	b	a
c	c	b	c	a	b	c	b	a	c	c	b	a	b	c	b	a	a

The 3-kernel contains $17 \leq 3^3$ sequences (also depends on the coding g)



		0	1	2	01	02	10	20	22	010	020	022	201	202	220	0220	2022
a	1	1	0	0	0	0	0	0	1	0	0	1	0	0	1	1	1
b	0	0	0	0	0	1	0	0	1	0	1	0	0	0	1	0	1
c	0	0	0	1	0	0	0	1	0	0	0	1	0	0	0	1	1

The 3-kernel contains $17 \leq 3^3$ sequences (also depends on the coding g , here 7 sequences: 011 is missing)

Recall that the Grundy function is unbounded (e.g., game of Nim)
How to generalize k -automatic sequences to an “infinite alphabet”?

DEFINITION [16]

$\mathbf{x} \in \mathbb{Z}^{\mathbb{N}}$ is **k -regular** if the \mathbb{Z} -module generated by $\text{Ker}_k(\mathbf{x})$ is finitely generated, i.e., there exists $\mathbf{t}_1, \dots, \mathbf{t}_\ell \in \mathbb{Z}^{\mathbb{N}}$ such that

$$\langle \text{Ker}_k(\mathbf{x}) \rangle = \langle \mathbf{t}_1, \dots, \mathbf{t}_\ell \rangle.$$

\mathbb{Z} is embedded in fields such as \mathbb{Q} , \mathbb{R} or \mathbb{C} . Thus the sequences can be seen as elements of $\mathbb{Q}^{\mathbb{N}}$ which is a \mathbb{Q} -vector space.

The orbit of \mathbf{x} under the action of the operators of decimation $\partial_{k,r}$ remains in a finite dimensional vector space.

ORIGINAL DEFINITION (ALLOUCHE–SHALLIT 1990)

Let R be a ring containing a commutative Noetherian ring R' .
A sequence \mathbf{x} in $R^{\mathbb{N}}$ is (R', k) -regular if there exists $\mathbf{t}_1, \dots, \mathbf{t}_\ell \in R^{\mathbb{N}}$ such that every sequence in $\text{Ker}_k(\mathbf{x})$ is an R' -linear combination of $\mathbf{t}_1, \dots, \mathbf{t}_\ell$.

$\langle \text{Ker}_k(\mathbf{x}) \rangle$ is a submodule of a finitely generated R' -module (in general, this does not imply that the submodule itself is finitely generated).

Since R' is assumed to be Noetherian, one can show that every submodule of a finitely generated R' -module is finitely generated and thus $\langle \text{Ker}_k(\mathbf{x}) \rangle$ is finitely generated.

EXAMPLE

The base-2 sum-of-digits function s_2 gives the sequence

$$(s_2(n))_{n \geq 0} = 0, 1, 1, 2, 1, 2, 2, 3, 1, 2, 2, 3, 2, 3, 3, 4, 1, 2, 2, \dots$$

Clearly this sequence is unbounded: $s_2(2^n - 1) = n$ for all n .

The \mathbb{Z} -module generated by its 2-kernel is generated¹ by the sequence $(s_2(n))_{n \geq 0}$ itself and the constant sequence $(1)_{n \geq 0}$:

$$s_2(2n) = s_2(n), \quad s_2(2n + 1) = s_2(n) + 1$$

PROPOSITION

Let s be a sequence taking finitely many different values. Let $k \geq 2$. The sequence is k -automatic if and only if it is k -regular.

Remark: intermediate notion of k -synchronized sequences. [17]

¹If it's unclear how to conclude from these two relations, wait a few slides, we will do it in details on another example with Nim.

With every k -regular sequence $(\mathbf{x}(n))_{n \geq 0} \in \mathbb{Z}^{\mathbb{N}}$ is associated a **linear representation** (λ, μ, ν) : there exist $r \in \mathbb{N}_{>0}$, $\lambda \in \mathbb{Z}^{1 \times r}$, $\nu \in \mathbb{Z}^{r \times 1}$ and a matrix-valued morphism $\mu : \{0, \dots, k-1\} \rightarrow \mathbb{Z}^{r \times r}$ such that

$$\mathbf{x}(n) = \lambda \mu(c_0 \cdots c_\ell) \nu$$

for all $c_\ell, \dots, c_0 \in \{0, \dots, k-1\}$ such that $\text{val}_k(c_\ell \cdots c_0) = n$.

The converse holds: if there exists a linear representation associated with canonical k -ary expansions (take into account the technicality of leading zeros), then the sequence is k -regular. See, Allouche–Shallit'03 Thm. 16.2.3 [7].

COROLLARY [16]

The n th term of a k -regular sequence can be computed with $\lfloor \log_k(n) \rfloor$ matrix multiplications.

If you are familiar with rational series:

THEOREM

A sequence $\mathbf{x}(n)$ is k -regular if and only if the formal series

$$\sum_{w \in \{0, \dots, k-1\}^*} \mathbf{x}(\text{val}_k(w)) w$$

is recognizable (in the sense of Berstel–Reutenauer [18]).

The sequence $s_2 = (s_2(n))_{n \geq 0}$ has a (base-2) linear representation given by

$$\lambda = \begin{pmatrix} 0 & 1 \end{pmatrix}, \mu(0) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \mu(1) = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \nu = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Think of $(s_2(n))$ and (1) as a basis. Matrix representation of the linear operators $\partial_{2,0}$ and $\partial_{2,1}$ in that basis:

$$\partial_{2,0}(s_2(n)) = (s_2(n)) \text{ and } \partial_{2,0}(1) = (1);$$

$$\partial_{2,1}(s_2(n)) = (s_2(n)) + (1) \text{ and } \partial_{2,1}(1) = (1).$$

SOME APPLICATIONS OF ITERATED MORPHISMS

- ▶ overlaps are avoidable over a 2-letter alphabet [19, 20, 21]

abbabaabbaababbabaababbaabbabaab ...

$$a \mapsto ab, \quad b \mapsto ba;$$

- ▶ cubes are avoidable over a 3-letter alphabet

abb|ab|a|abb|a|ab|abb|ab|a|ab|abb|a|abb|ab|a|ab ...

321312321231321 ...

$$3 \mapsto 321, \quad 2 \mapsto 31, \quad 1 \mapsto 2.$$

V. Keränen (ICALP'1992) avoiding abelian squares [22]

$a \mapsto abcacdc bcdcadcd b d a b a c a b a d b a b c b d b c b a c b c d c a c b a b d$
 $abacadc b c d c a c d b c b a c b c d c a c d c b d c d a d b d c b c a;$

$b \mapsto b c d b d a d c d a d b a d a c a b c b d b c b a c b c d c a c d c b d c d a d b d c b c a$
 $b c b d b a d c d a d b d a c d c b d c d a d b d a d c a d a b a c a d c d b;$

$c \mapsto c d a c a b a d a b a c b a b d b c d c a c d c b d c d a d b d a d c a d a b a c a d c d b$
 $c d c a c b a d a b a c a b d a d c a d a b a c a b a d b a b c b d b a d a c;$

$d \mapsto d a b d b c b a b c b d c b c a c d a d b d a d c a d a b a c a b a d b a b c b d b a d a c$
 $d a d b d c b a b c b d b c a b a d b a b c b d b c b a c b c d c a c b a b d;$

85-uniform morphism

J. Cassaigne, J. D. Currie, L. Schaeffer, J. Shallit, *Avoiding Three Consecutive Blocks of the Same Size and Same Sum* [23]

$$\varphi : 0 \mapsto 03, 1 \mapsto 43, 3 \mapsto 1, 4 \mapsto 01$$

$$\varphi^\omega(0) = 031430110343430310110110314303434303434 \dots$$

has no additive cube, e.g., 041340.

Prouhet's problem (1851) – Tarry – Escott [24, 25, 26]

Mémoire sur quelques relations entre les puissances de nombres

Partition $\{0, \dots, 2^n - 1\}$ such that

$$\{0, \dots, 2^n - 1\} = \{a_1, \dots, a_{2^{n-1}}\} \cup \{b_1, \dots, b_{2^{n-1}}\}$$

$$\sum_{j=1}^{2^{n-1}} a_j = \sum_{j=1}^{2^{n-1}} b_j, \quad \sum_{j=1}^{2^{n-1}} a_j^2 = \sum_{j=1}^{2^{n-1}} b_j^2, \quad \dots, \quad \sum_{j=1}^{2^{n-1}} a_j^{n-1} = \sum_{j=1}^{2^{n-1}} b_j^{n-1}$$

$$\prod_{i=0}^{\infty} (1 - X^{2^i}) = \sum_{j=0}^{\infty} t_j X^j$$

EXTENSION TO A MULTIDIMENSIONAL SETTING

There are natural extensions [10]

- Image under a coding of a fixed point of a k -uniform morphism;

$$a \mapsto \begin{array}{|c|c|} \hline b & a \\ \hline a & b \\ \hline \end{array} \quad b \mapsto \begin{array}{|c|c|} \hline a & b \\ \hline a & b \\ \hline \end{array}$$

- Sequence of outputs of a DFAO fed with base- k expansions;



$$q \xrightarrow{\begin{pmatrix} 0 \\ 1 \end{pmatrix}} q' \quad \text{padding shorter expansions with 0}$$

- Sets of indices defined by a first order formula in $\langle \mathbb{N}, +, V_k \rangle$;

$$\psi_a(x, y) \equiv \dots$$


- Finiteness of the k -kernel.

Remark: one could define (k, ℓ) -automatic sequences
(see, again, Allouche–Shallit 2003, [7, Chap. 14]).




O. Salon, Suites automatiques à multi-indices, *Séminaire de théorie des nombres*, Bordeaux, 1986–1987, exposé 4. [44]

Tracking the past




$$x(12, 10) \quad \text{rep}_2(12) = 1100, \quad \text{rep}_2(10) = 1010$$

Tracking the past




$$x(12, 10) \quad \text{rep}_2(12) = 1100, \quad \text{rep}_2(10) = 1010$$

Tracking the past




$$x(6, 5) \rightarrow x(12, 10) \quad \text{rep}_2(6) = 110, \quad \text{rep}_2(5) = 101$$

Tracking the past



$$x(3, 2) \rightarrow x(6, 5) \rightarrow x(12, 10) \quad \text{rep}_2(3) = 11, \quad \text{rep}_2(2) = 10$$

Tracking the past



$$x(1,1) \rightarrow x(3,2) \rightarrow x(6,5) \rightarrow x(12,10) \quad \text{rep}_2(3) = 1, \quad \text{rep}_2(2) = 1$$

Definition of the k -kernel in a multidimensional setting

DEFINITION

Consider a bi-dimensional sequence $\mathbf{x} = (x(m, n))_{m, n \geq 0}$.
It is a set of bi-dimensional subsequences:


$$\text{Ker}_k(\mathbf{x}) = \{ (x(k^i m + r, k^i n + s))_{m, n \geq 0} \mid i \geq 0, 0 \leq r, s < k^i \}.$$

This corresponds to selecting the suffixes


$$(0^{i-p} r_p \cdots r_1, 0^{i-q} s_q \cdots s_1)$$

where $\text{rep}_k(r) = r_p \cdots r_1$ and $\text{rep}_k(s) = s_q \cdots s_1$.


Some of these subsequences $(0, 0)$




Some of these subsequences $(1, 0)$




Some of these subsequences $(0, 1)$




Some of these subsequences (1, 1)




Some of these subsequences (00,00)



Some of these subsequences (01,00)



Some of these subsequences (10,00)



\rightsquigarrow We can define multidimensional k -regular sequences.

The \mathbb{Z} -module generated by $\text{Ker}_k(\mathbf{x})$ is finitely generated.

PROPOSITION (EXERCISE)

For the game of Nim, $(\mathcal{G}_N(m, n))_{m, n \geq 0}$ is 2-regular.

Proof. We have

$$\begin{aligned}\mathcal{G}_N(2m, 2n) &= 2m \oplus 2n &= 2\mathcal{G}_N(m, n) \\ \mathcal{G}_N(2m+1, 2n) &= (2m+1) \oplus 2n &= 2\mathcal{G}_N(m, n) + 1 \\ \mathcal{G}_N(2m, 2n+1) &= 2m \oplus (2n+1) &= 2\mathcal{G}_N(m, n) + 1 \\ \mathcal{G}_N(2m+1, 2n+1) &= (2m+1) \oplus (2n+1) &= 2\mathcal{G}_N(m, n)\end{aligned}$$

thus the 2-kernel is generated by $(\mathcal{G}_N(m, n))_{m, n \geq 0}$ and the constant sequence (1). □

Is that clear for any element of the 2-kernel?

Can $(\mathcal{G}_N(8m+5, 8n+2))_{m,n \geq 0}$ be expressed as a \mathbb{Z} -linear combination of these two sequences?

$$\begin{aligned}\mathcal{G}_N(8m+5, 8n+2) &= \mathcal{G}_N(2(4m+2)+1, 2(4n+1)) \\ &= 2\mathcal{G}_N(4m+2, 4n+1) + 1 \\ &= 2\mathcal{G}_N(2(2m+1), 2(2n+1)) + 1 \\ &= 2[2\mathcal{G}_N(2m+1, 2n) + 1] + 1 \\ &= 4\mathcal{G}_N(2m+1, 2n) + 3 \\ &= 4[2\mathcal{G}_N(m, n) + 1] + 3 \\ &= 8\mathcal{G}_N(m, n) + 7.\end{aligned}$$

Meaning of these relations within the table:

9	8	11	10	13	12	15	14	1	0
8	9	10	11	12	13	14	15	0	1
7	6	5	4	3	2	1	0	15	14
6	7	4	5	2	3	0	1	14	15
5	4	7	6	1	0	3	2	13	12
4	5	6	7	0	1	2	3	12	13
3	2	1	0	7	6	5	4	11	10
2	3	0	1	6	7	4	5	10	11
1	0	3	2	5	4	7	6	9	8
0	1	2	3	4	5	6	7	8	9

First few values of $\mathcal{G}_N(m, n)$.

$$\mathcal{G}_N(m, n) \mapsto \begin{array}{|c|c|} \hline 2\mathcal{G}_N(m, n) + 1 & 2\mathcal{G}_N(m, n) \\ \hline 2\mathcal{G}_N(m, n) & 2\mathcal{G}_N(m, n) + 1 \\ \hline \end{array}$$

For the game of Wythoff, first few values of $(x, y) \mapsto \mathcal{G}_W(x, y)$


\vdots											\ddots
9	9	10	11	12	8	7	13	14	15	16	
8	8	6	7	10	1	2	5	3	4	15	
7	7	8	6	9	0	1	4	5	3	14	
6	6	7	8	1	9	10	3	4	5	13	
5	5	3	4	0	6	8	10	1	2	7	
4	4	5	3	2	7	6	9	0	1	8	
3	3	4	5	6	2	0	1	9	10	12	
2	2	0	1	5	3	4	8	6	7	11	
1	1	2	0	4	5	3	7	8	6	10	
0	0	1	2	3	4	5	6	7	8	9	...
	0	1	2	3	4	5	6	7	8	9	...

Not so many results are known

- ▶ U. Blass, A.S. Fraenkel, The Sprague-Grundy function for Wythoff's game. Theoret. Comput. Sci. 75 (1990), no. 3, 311–333. [27]
- ▶ Y. Jiao, On the Sprague-Grundy values of the \mathcal{F} -Wythoff game. Electron. J. Combin. 20 (2013). [28]
- ▶ A. Gu, Sprague-Grundy values of the \mathcal{R} -Wythoff game. Electron. J. Combin. 22 (2015). [29]
- ▶ M. Weinstein, Invariance of the Sprague-Grundy function for variants of Wythoff's game. Integers 16 (2016). [30]

It's challenging, we quote the book [1, A. Siegel, p. 200]:

“No general formula is known for computing arbitrary \mathcal{G} -values of WYTHOFF. In general, they appear chaotic, though they exhibit a striking fractal-like pattern ... Despite this apparent chaos, the \mathcal{G} -values nonetheless have a high degree of geometric regularity.”



$$\mathcal{G}_W(m, n), \quad m, n \leq 100$$

Defining the MeX function for a list

```
mex[{}] = 0;  
mex[l_] := Min[Complement[Table[i, {i, 0, Max[l] + 1}], l]]
```

List the available pairs of positions

```
In[19]:= pos[x_ /; x > 0, y_ /; y > 0] := Union[Table[{i, y}, {i, 0, x - 1}],  
  Table[{x, i}, {i, 0, y - 1}], Table[{x - i, y - i}, {i, 1, Min[{x, y}]}]];  
pos[0, y_ /; y > 0] := Table[{0, i}, {i, 0, y - 1}];  
pos[x_ /; x > 0, 0] := Table[{i, 0}, {i, 0, x - 1}];
```

Compute the Sprague-Grundy function for Wythoff's game

```
In[28]:= w[{0, 0}] = 0;  
w[{x_, y_}] := w[{x, y}] = mex[Map[w[#] &, pos[x, y]]]
```

```
In[50]:= Timing[  
  t = Table[w[{i, j}], {i, 0, 100}, {j, 0, 100}];]
```

```
Out[50]=  
  
{2.03782, Null}
```

Give nice RGB colors, 0 is mapped to Red, other values from Green to Blue

```
In[51]:= Graphics[Raster[Map[If[# > 0, {0, Max[t] - #, #} / Max[t], {1, 0, 0}] &, t, {2}]]]
```

Some results from Blass and Fraenkel [27]:

- ▶ On every parallel to the main diagonal, $\mathcal{G}_W(n, n + j)$ takes every possible value.
- ▶ Points with Grundy value 1 are “close” to those with value 0.
- ▶ Recursive algorithm to determine the points with Grundy value 1.

Additive periodicity of the rows of $(\mathcal{G}_W(m, n))$


DEFINITION

A sequence $(a_j)_{j \geq 0}$ is *additively periodic* if

$$\exists p, q, \forall j \geq q : a_{j+p} = a_j + p.$$

- ▶ A. Dress, A. Flammenkamp and N. Pink, Additive periodicity of the Sprague–Grundy function of certain Nim games, Adv. in Appl. Math. 22, 249–270 (1999). [31]
- ▶ H. A. Landman, A simple FSM²-based proof of the additive periodicity of the Sprague-Grundy function of Wythoff's game, More Games of No Chance, 2002. [32]

As an example, the row $\mathcal{G}_W(5, n)$ is such that
 for all $n \geq 27$, $\mathcal{G}_W(5, n + 24) = \mathcal{G}_W(5, n) + 24$



graph of $\mathcal{G}_W(5, n) - n$ for $50 \leq n \leq 150$.

Observe that the Fibonacci word is coding the \mathcal{P} -positions:

We have a Beatty sequence $\frac{1}{\varphi} + \frac{1}{\varphi^2} = 1$,

$(\lfloor n\varphi \rfloor)_{n \geq 1}$ and $(\lfloor n\varphi^2 \rfloor)_{n \geq 1}$ make a partition of $\mathbb{N}_{>0}$ [33]


1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, ...

This corresponds to the Sturmian (Fibonacci) word:

$\mathbf{f} = abaababaab \dots = 2122121221 \dots$

$$\mathbf{f}_n = \lfloor \varphi(n+2) \rfloor - \lfloor \varphi(n+1) \rfloor, \quad \forall n \geq 0.$$

A word is Sturmian iff
it is irrational mechanical [34].



PROPOSITION


The n th \mathcal{P} -position of Wythoff's game is given by the positions (starting with 1) of the n th a and n th b in \mathbf{f} .

Link with (Zeckendorf) numeration system [35]:

The Fibonacci word is the fixed point of $a \mapsto ab$, $b \mapsto a$.

What could be the analogue to

- ▶ Image under a coding of a fixed point of a k -uniform morphism;
- ▶ Sequence of outputs of a DFAO fed with $\text{base-}k$ expansions;



$\varepsilon, 1, 10, 100, 101, 1000, 1001, 1010, 10000, 10001, 10010, 10100, \dots$

This is exactly the language of (greedy) Fibonacci representations $\text{rep}_F(\mathbb{N})$. We have a “**Fibonacci-automatic sequence**”.

We have a general result (indexing **starts with 0**):

GENERAL THEOREM “MORPHIC \Rightarrow AUTOMATIC” [40]

Let A be an ordered alphabet. Let $\mathbf{w} \in A^{\mathbb{N}}$ be an infinite word, fixed point $f^{\omega}(a)$ of a morphism $f : A^* \rightarrow A^*$.

- ▶ associate with f a DFA \mathcal{M} over the alphabet $\{0, \dots, \max |f(b)| - 1\}$;
- ▶ A is the set of states;
- ▶ the initial state is a , all states are final;
- ▶ if $f(b) = c_0 \cdots c_m$, then $b \xrightarrow{j} c_j$, $j \leq m$;
- ▶ consider the language L accepted by \mathcal{M} except words starting with 0;
- ▶ genealogically order L : $L = \{w_0 < w_1 < w_2 < \cdots\}$.

The n th symbol of \mathbf{w} , $n \geq 0$, is $\boxed{\mathcal{M} \cdot w_n}$.

\rightsquigarrow The ordered language L plays the rôle of the base- k numeration system. We have a “ **L -automatic sequence**”.

Many links with non-standard numeration systems

J. Shallit (1988), J.-P. Allouche, E. Cateland, et al. (1997), J.-P. Allouche, K. Scheicher, R. Tichy (2000), M. R. (2000),... [36, 37, 38, 39, 40]

Starting with index 0: $f_0 f_1 f_2 \cdots = abaab \cdots$

$f_j = a$ iff $\text{rep}_F(j)$ ends with 0; $f_j = b$ iff $\text{rep}_F(j)$ ends with 1.

We can still “keep track of the past”: since $a \mapsto ab$, $b \mapsto a$, the n th symbol b comes from the n th a in \mathbf{f} .

ε	1	10	100	101	1000	1001	1010	10000	10001	10010	10100	10101	
a	b	a	a	b	a	b	a	a	b	a	a	b	\cdots

$f_j = a$ and $\text{rep}_F(j) = u0$
iff

$f_{\ell-1} = a, f_{\ell} = b$ with $\ell = \text{val}_F(u01)$.

$f_j = b$ and $\text{rep}_F(j) = v1$ iff $f_{\ell} = a$ with $\ell = \text{val}_F(v10)$.

Putting together the four previous slides:

PROPOSITION

The n th \mathcal{P} -position of Wythoff's game is given by the positions (starting with 1) of the n th a and n th b in \mathbf{f} .

(x, y) , with $x < y$, is a \mathcal{P} -position iff

$$(\text{rep}_F(x-1), \text{rep}_F(y-1)) = (u0, u01)$$

where u is a greedy (valid) F -representation.

What can be said about the form of $\text{rep}_F(x)$ and $\text{rep}_F(y)$?

$$\text{rep}_F(x) = \text{rep}_F(\text{val}_F(u0) + 1)$$

$$\text{rep}_F(y) = \text{rep}_F(\text{val}_F(u01) + 1)$$

- First case³ : $u = u'0$

$$\text{rep}_F(x) = \text{rep}_F(\text{val}_F(u'00) + 1) = u'01 \text{ ends with no zero}$$

$$\text{rep}_F(y) = \text{rep}_F(\text{val}_F(u'001) + 1) = u'010 \text{ shift by one zero}$$


- Second case : $u = u'1$ (u' ends with 0)

$$\text{rep}_F(x) = \text{rep}_F(\text{val}_F(u'10) + 1) = u'\mathbf{11}$$

$$\text{rep}_F(y) = \text{rep}_F(\text{val}_F(u'101) + 1) = u'\mathbf{110}$$

Normalize $u'\mathbf{11}$ / $u'\mathbf{110}$ or compute the successor of $u'10$ / $u'101$
Transducer (R to L) computing the successor, see [41, Frougny'97].

³The first values may be checked by hand.



$$s \xleftarrow{\begin{pmatrix} x1 \\ x1 \end{pmatrix}} s \xleftarrow{\begin{pmatrix} 0 \\ 01 \end{pmatrix}} r \left[\xleftarrow{\begin{pmatrix} 0 \\ 0 \end{pmatrix}} q \xleftarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} r \right] \xleftarrow{\begin{pmatrix} 0 \\ 0 \end{pmatrix}} q \xleftarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} r \xleftarrow{\begin{pmatrix} 0 \\ . \end{pmatrix}} p \text{ :from right}$$

$$\underbrace{x10(01)^n 010}_{u'} \rightarrow x101(00)^n 00 \quad 2n + 2 \text{ zeroes}, n \geq 0$$

$$\underbrace{x10(01)^n 0101}_{u'} \rightarrow x101(00)^n 000 \quad \text{shift by one zero}, n \geq 0$$

$$\underbrace{1(01)^n 010}_{u'} \rightarrow 100(00)^n 00 \quad 2n + 4 \text{ zeroes}, n \geq 0$$

$$\underbrace{1(01)^n 0101}_{u'} \rightarrow 100(00)^n 000 \quad \text{shift by one zero}, n \geq 0$$

COROLLARY (A. S. FRAENKEL, 1982 [42])

(x, y) , with $x < y$, is a \mathcal{P} -position iff $\text{rep}_F(x)$ ends with an even number of zeroes and $\text{rep}_F(y) = \text{rep}_F(x)0$.

SHAPE-SYMMETRY

Question: *What can be said about the (morphic) structure of the \mathcal{P} -positions of Wythoff's ♔ game?* [43]

$$(P_{i,j})_{i,j \geq 0} = \begin{array}{cccccccccccc|c} & \vdots & & & & & & & & & & & \ddots \\ 0 & 0 & 0 & 0 & 0 & 0 & \mathbf{1} & 0 & 0 & 0 & 0 & & \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & & \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & & \\ 0 & 0 & 0 & 0 & \mathbf{1} & 0 & 0 & 0 & 0 & 0 & 0 & & \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \mathbf{1} & & \\ 0 & 0 & 0 & \mathbf{1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & & \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \mathbf{1} & 0 & 0 & 0 & & \\ 0 & 0 & 0 & 0 & 0 & \mathbf{1} & 0 & 0 & 0 & 0 & 0 & & \\ 0 & \mathbf{1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & & \\ 0 & 0 & \mathbf{1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & & \\ \mathbf{1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \end{array}$$

Let's try something...

$$\varphi_W : a \mapsto \begin{array}{|c|c|} \hline c & d \\ \hline a & b \\ \hline \end{array} \quad b \mapsto \begin{array}{|c|} \hline e \\ \hline i \\ \hline \end{array} \quad c \mapsto \begin{array}{|c|c|} \hline i & j \\ \hline \end{array} \quad d \mapsto \begin{array}{|c|} \hline i \\ \hline \end{array} \quad e \mapsto \begin{array}{|c|c|} \hline f & b \\ \hline \end{array}$$

$$f \mapsto \begin{array}{|c|c|} \hline h & d \\ \hline g & b \\ \hline \end{array} \quad g \mapsto \begin{array}{|c|c|} \hline h & d \\ \hline f & b \\ \hline \end{array} \quad h \mapsto \begin{array}{|c|c|} \hline i & m \\ \hline \end{array} \quad i \mapsto \begin{array}{|c|c|} \hline h & d \\ \hline i & m \\ \hline \end{array}$$

$$j \mapsto \begin{array}{|c|} \hline c \\ \hline k \\ \hline \end{array} \quad k \mapsto \begin{array}{|c|c|} \hline c & d \\ \hline l & m \\ \hline \end{array} \quad l \mapsto \begin{array}{|c|c|} \hline c & d \\ \hline k & m \\ \hline \end{array} \quad m \mapsto \begin{array}{|c|} \hline h \\ \hline i \\ \hline \end{array}$$


and the coding

$$\mu_W : a, e, g, j, l \mapsto 1, \quad b, c, d, f, h, i, k, m \mapsto 0$$

The idea:

SHAPE-SYMMETRIC MORPHISM (A. MAES, 1999 [47])

If P is the infinite bidimensional picture that is the fixed point of φ , then for all $i, j \in \mathbb{N}$, if $\varphi(P_{i,j})$ is a block of size $k \times \ell$ then $\varphi(P_{j,i})$ is of size $\ell \times k$



The details: Let $d \geq 2$

A d -dimensional picture over A is a map

$$x : \llbracket 0, s_1 - 1 \rrbracket \times \cdots \times \llbracket 0, s_d - 1 \rrbracket \rightarrow A$$

(s_1, \dots, s_d) is the **shape** of x ; if $s_i < \infty$, for all i , x is **bounded**.

The set of bounded pictures over A is denoted by $\mathcal{B}_d(A)$.

If for some $i \in \llbracket 1, d \rrbracket$, $|x|_{\hat{i}} = |y|_{\hat{i}} = (s_1, \dots, s_{i-1}, s_{i+1}, \dots, s_d)$, then we define the **concatenation** of x and y *in the direction i* to be the d -dimensional picture $x \odot^i y$ of shape

$$(s_1, \dots, s_{i-1}, |x|_i + |y|_i, s_{i+1}, \dots, s_d).$$

$$x = \begin{array}{|c|c|} \hline a & b \\ \hline c & d \\ \hline \end{array} \quad \text{and} \quad y = \begin{array}{|c|c|c|} \hline a & a & b \\ \hline b & c & d \\ \hline \end{array}$$

of shape respectively $|x| = (2, 2)$ and $|y| = (2, 3)$.

Since $|x|_{\hat{2}} = |y|_{\hat{2}} = 2$, we get

$$x \odot^2 y = \begin{array}{|c|c|c|c|c|} \hline a & b & a & a & b \\ \hline c & d & b & c & d \\ \hline \end{array}.$$

However $x \odot^1 y$ is not defined because $2 = |x|_{\hat{1}} \neq |y|_{\hat{1}} = 3$.

A map $\gamma: A \rightarrow \mathcal{B}_d(A)$ cannot necessarily be extended to a morphism $\gamma: \mathcal{B}_d(A) \rightarrow \mathcal{B}_d(A)$.

$$\gamma: a \mapsto \begin{bmatrix} b & d \\ a & a \end{bmatrix}, \quad b \mapsto \begin{bmatrix} b \\ c \end{bmatrix}, \quad c \mapsto \begin{bmatrix} a & a \end{bmatrix}, \quad d \mapsto \begin{bmatrix} d \end{bmatrix}.$$

$$\odot^2: |\gamma(c)|_2 = |\gamma(d)|_2 = 1, \quad |\gamma(a)|_2 = |\gamma(b)|_2 = 2,$$

$$\odot^1: |\gamma(a)|_1 = |\gamma(c)|_1 = 2, \quad |\gamma(d)|_1 = |\gamma(b)|_1 = 1.$$

$$x = \begin{bmatrix} c & d \\ a & b \end{bmatrix}, \quad \gamma(x) = \begin{bmatrix} a & a & d \\ b & d & b \\ a & a & c \end{bmatrix}$$

but $\gamma^2(x)$ is not well-defined!

DEFINITION

If for all $a \in A$ and all $n \geq 1$, $\gamma^n(a)$ is well-defined from $\gamma^{n-1}(a)$, then γ is said to be a **d -dimensional morphism**.

\rightsquigarrow the images of any two symbols on a row (resp. column) have the same number of rows (resp. columns).

DEFINITION

Let $\gamma: \mathcal{B}_d(A) \rightarrow \mathcal{B}_d(A)$ be a d -dimensional morphism having the d -dimensional infinite word x as a fixed point.


This word is **shape-symmetric with respect to γ** if, for all permutations ν of $\llbracket 1, d \rrbracket$, we have, for all $n_1, \dots, n_d \geq 0$,

$$|\gamma(x(n_1, \dots, n_d))| = (s_1, \dots, s_d)$$

$$\Downarrow$$

$$|\gamma(x(n_{\nu(1)}, \dots, n_{\nu(d)}))| = (s_{\nu(1)}, \dots, s_{\nu(d)}).$$


Reconsider our map φ (one can indeed prove that it is a d -dimensional morphism having a shape-symmetric fixed point).



sizes : 1, 2, 3, 5

\dots	\mapsto	<i>i</i>	<i>m</i>	<i>i</i>	<i>i</i>	j	<i>i</i>	<i>m</i>	<i>i</i>	\mapsto	\dots
		<i>h</i>	<i>d</i>	<i>h</i>	<i>c</i>	<i>d</i>	<i>h</i>	<i>d</i>	<i>h</i>		
		<i>i</i>	<i>m</i>	<i>i</i>	l	<i>m</i>	<i>i</i>	<i>m</i>	<i>i</i>		
		<i>h</i>	<i>d</i>	<i>c</i>	<i>h</i>	<i>d</i>	<i>h</i>	<i>d</i>	e		
		<i>i</i>	<i>m</i>	<i>k</i>	<i>i</i>	<i>m</i>	g	<i>b</i>	<i>i</i>		
		<i>i</i>	j	<i>i</i>	<i>f</i>	<i>b</i>	<i>i</i>	<i>m</i>	<i>i</i>		
		<i>c</i>	<i>d</i>	e	<i>h</i>	<i>d</i>	<i>h</i>	<i>d</i>	<i>h</i>		
		a	<i>b</i>	<i>i</i>	<i>i</i>	<i>m</i>	<i>i</i>	<i>m</i>	<i>i</i>		

size : 8,...



Before proceeding to the proof (of morphic structure of the \mathcal{P} -positions of Wythoff)... How did we get that?

\vdots								\ddots
0 0	0	0 0	0 1	0	0 0	0		
0 0	0	0 0	0 0	0	0 0	0		
0 0	0	0 0	0 0	0	0 0	0		
0 0	0	0 1	0 0	0	0 0	0		
0 0	0	0 0	0 0	0	0 0	1		
0 0	0	1 0	0 0	0	0 0	0		
0 0	0	0 0	0 0	0	0 0	0		
0 0	0	0 0	1 0	0	0 0	0		
0 1	0	0 0	0 0	0	0 0	0		
0 0	1	0 0	0 0	0	0 0	0		
1 0	0	0 0	0 0	0	0 0	0	\dots	

shape sequence: 2122121221...

THEOREM [43]

The morphism φ_W and the coding μ_W give the 2-dimensional infinite word coding the \mathcal{P} -positions of Wythoff.

Proof. We can do the same⁴ as for the uni-dimensional case:
We associate with φ an automaton with input alphabet

$$\left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$$


$$\varphi(r) = \begin{array}{|c|c|} \hline u & v \\ \hline s & t \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline s & t \\ \hline \end{array}, \quad \begin{array}{|c|} \hline u \\ \hline s \\ \hline \end{array} \quad \text{or} \quad \begin{array}{|c|} \hline s \\ \hline \end{array}$$

we have transitions like

$$r \xrightarrow{\begin{pmatrix} 0 \\ 0 \end{pmatrix}} s, \quad r \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} t, \quad r \xrightarrow{\begin{pmatrix} 0 \\ 1 \end{pmatrix}} u, \quad r \xrightarrow{\begin{pmatrix} 1 \\ 1 \end{pmatrix}} v.$$

⁴Similar to the general theorem: the n th symbol is obtained by feeding a DFA with the n th word of the accepted language.

From morphism to automaton, we get



1) If all states are assumed to be final, this automaton accepts the words

$$\begin{pmatrix} u \\ v \end{pmatrix}$$

where $|u| = |v|$ and u, v are both valid F -representation (possibly padded with zeroes).


2) If we restrict to the “blue” part, this automaton accepts the words

$$\begin{pmatrix} 0w_1 \cdots w_\ell \\ w_1 \cdots w_\ell 0 \end{pmatrix} \text{ and } \begin{pmatrix} w_1 \cdots w_\ell 0 \\ 0w_1 \cdots w_\ell \end{pmatrix}$$

where $w_1 \cdots w_\ell$ is a valid F -representation.


3) Now, if the set of final states is $\{a, e, g, j, l\}$, we have the extra condition that $w_1 \cdots w_\ell$ **ends with an even number of zeroes**.

With our previous characterization of \mathcal{P} -positions, this concludes the proof. □



Initial blocks of some 3-dimensional shape-symmetric picture
[47, Maes' thesis p. 107].

ASSOCIATED DECISION PROBLEMS

 **UMH**
UNIVERSITÉ DE MONT-HELIAN
Faculté des Sciences
Institut de Mathématique
et d'Informatique
Service de Logique Mathématique et Algèbre

MORPHIC PREDICATES AND APPLICATIONS TO THE DECIDABILITY OF ARITHMETIC THEORIES

Arnaud MAES

Dissertation originale présentée
pour l'obtention du grade académique de
DOCTEUR EN SCIENCES

Janvier 1999

Some Results from Maes' papers:

- ▶ Determining whether or not a map $\mu: \mathcal{B}_d(A) \rightarrow \mathcal{B}_d(A)$ is a d -dimensional morphism is a **decidable** problem.
- ▶ If μ is prolongable on a letter a , then it is **decidable** whether or not the fixed point $\mu^\omega(a)$ is shape-symmetric.

[45, 46, 47]

ADDING OR REMOVING MOVES

↪ How can we alter the **set of moves**  to keep the same set of \mathcal{P} -positions?

REMARK

This means that several rule-sets (different “games”) could lead to the same set of \mathcal{P} -positions.

In a subtraction game, observe that a move can be adjoined (*without altering the set of \mathcal{P} -positions*) if and only if

it does not belong to $\mathcal{P} - \mathcal{P}$.

COROLLARY

We can adjoin the move $(i, j)_{i < j}$ to Wythoff's rule-set iff

$$(i, j) \neq ([n\varphi] - [m\varphi], [n\varphi^2] - [m\varphi^2]) \quad \forall n > m \geq 0$$

$$\text{and } (i, j) \neq ([n\varphi] - [m\varphi^2], [n\varphi^2] - [m\varphi]) \quad \forall n > m \geq 0.$$

THEOREM (E. DUCHÊNE ET AL. 2010 [43])


$(i, j)_{i < j}$ may be adjoined iff there exist valid F -representations u, u' such that one the three properties is satisfied :

- ▶ $(\text{rep}_F(i - 1), \text{rep}_F(j - 1)) = (u0, u01)$
- ▶ $(\text{rep}_F(i - 2), \text{rep}_F(j - 2)) = (u0, u01)$
- ▶ $(\text{rep}_F(j - \lfloor i\varphi \rfloor - 2), \text{rep}_F(j - \lfloor i\varphi \rfloor - 2 + i)) = (u1, u'0);$

0	0	0	1	0	0	1	0	0	0	0
0	1	0	0	1	0	0	0	0	0	0
0	0	0	0	0	1	0	0	0	0	0
0	1	0	0	1	0	0	0	0	0	0
0	0	1	0	1	0	0	0	0	0	1
0	0	0	1	0	0	0	0	1	0	0
0	1	0	0	0	0	1	1	0	1	0
0	0	1	0	0	1	0	0	0	0	1
0	1	0	1	0	0	1	0	0	0	0
0	0	1	0	1	0	0	1	0	1	0
0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0

- There is no redundant move in Wythoff's game.

Question: Does the infinite 2-dimensional word on the previous slide have a shape-symmetric morphic structure?



We conjectured a morphism over 26 letters.

Another direction leads to the concept of *invariant games* [48, 49].

A game $G : \mathbb{N}^n \rightarrow 2^{\mathbb{N}^n}$ (assigning each position to a set of available moves) is **invariant** if there exists a set $I \subseteq \mathbb{N}^n$ such that, for all positions \mathbf{p} , we have

$$G(\mathbf{p}) = I \cap \{\mathbf{m} \in \mathbb{N}^n \mid \mathbf{m} \leq \mathbf{p}\}.$$

Otherwise stated, we may apply exactly the same moves to every position, with the only restriction that there are enough tokens left.

EXAMPLE

The game of Nim is invariant:

$$I_{\text{NIM}} = \{(i, 0) \mid i \geq 1\} \cup \{(0, j) \mid j \geq 1\}.$$

Wythoff's game is invariant:

$$I_{\text{WYTHOFF}} = I_{\text{NIM}} \cup \{(k, k) \mid k \geq 1\}.$$

For an example of non-invariant game, consider the following map,

$$G_{\text{EVEN}} : \mathbb{N}^2 \rightarrow 2^{\mathbb{N}^2},$$

$$(x, y) \mapsto \begin{cases} \{(i, 0) \mid i \in \llbracket 1, x \rrbracket\}, & \text{if } x + y \text{ is even;} \\ \{(i, i) \mid i \in \llbracket 1, \min\{x, y\} \rrbracket\}, & \text{otherwise.} \end{cases}$$

LINK WITH ANS

Actually, we have already seen *Abstract Numeration Systems*. . .

DEFINITION [50]

An **abstract numeration system** $\mathcal{S} = (L, A, <)$ is a regular language L over a totally ordered finite alphabet $(A, <)$.

- ▶ Enumerating the words in L using genealogical ordering provides a **one-to-one correspondence** between \mathbb{N} and L :


$$\text{rep}_{\mathcal{S}} : \mathbb{N} \rightarrow L, \quad \text{val}_{\mathcal{S}} : L \rightarrow \mathbb{N}.$$

- ▶ This generalizes any positional system U for which $\text{rep}_U(\mathbb{N})$ is regular.

P. Lecomte, M.R., Numeration systems on a regular language, *Theory Comput. Syst.* **34** (2001), 27–44.

ABSTRACT NUMERATION SYSTEMS


Example : consider a prefix-closed language $L = \{b, \varepsilon\}\{a, ab\}^*$



n	$\text{rep}_{\mathcal{S}}(n)$
0	ε
1	a
2	b
3	aa
4	ab
5	ba
6	aaa
7	aab
8	aba
9	baa
10	bab

ABSTRACT NUMERATION SYSTEMS

A non-positional ANS $L = a^*b^*$
#b




n	$\text{rep}_{\mathcal{S}}(n)$
0	ε
1	a
2	b
3	aa
4	ab
5	bb
6	aaa
7	aab
8	abb
9	bbb
10	$aaaa$

#a

ABSTRACT NUMERATION SYSTEMS

A non-positional ANS $L = a^*b^*$
#b




n	$\text{rep}_{\mathcal{S}}(n)$
0	ε
1	a
2	b
3	aa
4	ab
5	bb
6	aaa
7	aab
8	abb
9	bbb
10	$aaaa$

#a

ABSTRACT NUMERATION SYSTEMS

A non-positional ANS $L = a^*b^*$
#b




n	$\text{rep}_{\mathcal{S}}(n)$
0	ε
1	a
2	b
3	aa
4	ab
5	bb
6	aaa
7	aab
8	abb
9	bbb
10	$aaaa$

#a

ABSTRACT NUMERATION SYSTEMS

A non-positional ANS $L = a^*b^*$
#b




n	$\text{rep}_{\mathcal{S}}(n)$
0	ε
1	a
2	b
3	aa
4	ab
5	bb
6	aaa
7	aab
8	abb
9	bbb
10	$aaaa$

#a

ABSTRACT NUMERATION SYSTEMS

A non-positional ANS $L = a^*b^*$
#b




n	$\text{rep}_{\mathcal{S}}(n)$
0	ε
1	a
2	b
3	aa
4	ab
5	bb
6	aaa
7	aab
8	abb
9	bbb
10	$aaaa$

#a

ABSTRACT NUMERATION SYSTEMS

A non-positional ANS $L = a^*b^*$
#b




n	$\text{rep}_{\mathcal{S}}(n)$
0	ε
1	a
2	b
3	aa
4	ab
5	bb
6	aaa
7	aab
8	abb
9	bbb
10	$aaaa$

#a

ABSTRACT NUMERATION SYSTEMS

A non-positional ANS $L = a^*b^*$
#b




n	$\text{rep}_{\mathcal{S}}(n)$
0	ε
1	a
2	b
3	aa
4	ab
5	bb
6	aaa
7	aab
8	abb
9	bbb
10	$aaaa$

#a

ABSTRACT NUMERATION SYSTEMS

A non-positional ANS $L = a^*b^*$
#b



n	$\text{rep}_{\mathcal{S}}(n)$
0	ε
1	a
2	b
3	aa
4	ab
5	bb
6	aaa
7	aab
8	abb
9	bbb
10	$aaaa$

ABSTRACT NUMERATION SYSTEMS

A non-positional ANS $L = a^*b^*$

$$\text{val}_S(a^p b^q) = \frac{1}{2}(p+q)(p+q+1) + q = \binom{p+q+1}{2} + \binom{q}{1}$$

ε	a	b	aa	ab	bb	aaa	\dots
0	1	2	3	4	5	6	\dots

$$U_0 = 1, \quad U_1 = 2, \quad p(a) = 1, p(b) = 2$$

$$\text{Generalization : } \text{val}_\ell(a_1^{n_1} \cdots a_\ell^{n_\ell}) = \sum_{i=1}^{\ell} \binom{n_i + \cdots + n_\ell + \ell - i}{\ell - i + 1}.$$

$$\boxed{\forall n \in \mathbb{N}, \exists z_1, \dots, z_\ell : n = \binom{z_\ell}{\ell} + \binom{z_{\ell-1}}{\ell-1} + \cdots + \binom{z_1}{1}}$$


with the condition $z_\ell > z_{\ell-1} > \cdots > z_1 \geq 0$

There was already some form of abstract numeration system in Maes' Ph.D. thesis (1999) [47].

Rem. 6.9, p. 134, "*The set of codes of \mathbb{N} given by the above automaton is of course a regular language ... The language read by A is 0^*L . However, the above coding is not a numeration system in the sense of [6]. Indeed, the representation of a natural number is not obtained using a 'Euclidian division' algorithm.*"

\mathcal{S} -AUTOMATIC SEQUENCES


Two ingredients: an ANS $S = (a^*b^*, a < b)$ and a DFAO \mathcal{M} [39]



$$x_n = \mathcal{M} \cdot \text{rep}_S(n)$$

$\mathbf{x} = 01023031200231010123023031203120231002310123010123 \dots$

$S = (a^*b^*, a < b)$ padding symbol: #



\vdots
 220200200020000
 112222222222222
 020200200020000
 200000000000000
 022222222222222
 220200200020000
 112122122212222
 001011011101111...

THEOREM (A. MAES, M.R. [40])

An infinite word is morphic if and only if it is S -automatic for some abstract numeration system S .

\Rightarrow Already proven.

\Leftarrow We need to get rid off erasing morphisms.

Simulate the product of the two automata.

THEOREM (É. CHARLIER, T. KÄRKI, M.R. [54])

Let $d \geq 1$. The d -dimensional infinite word x is S -automatic, for some abstract numeration system $S = (L, \Sigma, <)$ where $\varepsilon \in L$, if and only if x is the image by a coding of a shape-symmetric infinite d -dimensional word.

SUMMARY

k -automatic sequence



k -uniform morphism

+ coding

[A. Cobham'72] [8]

\mathcal{S} -automatic sequence



non-erasing morphism

+ coding

[A. Maes, M.R.'02] [39, 40]

multidimensional setup

$$x : \mathbb{N}^d \rightarrow A$$

k -automatic sequence



morphism $g : A \rightarrow (A^k)^d$

+ coding

[O. Salon'87] [44]

\mathcal{S} -automatic sequence



“shape-symmetric” morphism

+ coding

[É. Charlier, T. Kärki, M.R.'09] [54]

GAMES WITH A FINITE SET OF MOVES

Games like Nim or Wythoff have an infinite set of moves.

IN ONE DIMENSION [1]

Every (invariant) finite subtraction game on one pile, *i.e.*, $I \subset \mathbb{N}$ is finite, has an ultimately periodic Grundy function.

Proof:

Let $m = \#I$ (max. number of options), then $\mathcal{G}(n) \leq m$ for all n .

Let $k = \max I$, there are $(m+1)^k$ possible k -tuples taking values in $\{0, \dots, m\}$. $\mathcal{G}(n)$ depends only on $\mathcal{G}(n-i)$ for $1 \leq i \leq k$.

Hence, there exist $i < j$

$$\mathcal{G}(i+n) = \mathcal{G}(j+n) \text{ for all } n \in \{0, \dots, k-1\}.$$

Thus $j-i$ is a period of \mathcal{G} with preperiod i . □

Another similar result

[1, A. SIEGEL, P. 188]

Consider an (invariant) finite subtraction game on one pile, with $I \subset \mathbb{N}$ as set of moves. If there exist $N \geq 0$ and $p \geq 1$ such that

$$\mathcal{G}(n + p) = \mathcal{G}(n), \quad \forall N \leq n < N + \max I$$

then $\mathcal{G}(n + p) = \mathcal{G}(n)$ for all $n \geq N$.

If we may **optionally split** a pile...

Definition from Wikipedia:

An **octal game** is played with tokens divided into heaps.

Two players take turns moving until no moves are possible.

Every move consists of selecting just one of the heaps, and either

- ▶ removing **all of the tokens in the heap**, leaving no heap,
- ▶ removing some but not all of the tokens, leaving **one smaller heap**, or
- ▶ removing some of the tokens and **dividing** the remaining tokens into **two nonempty heaps**.

Heaps other than the selected heap remain unchanged.

The last player to move wins in normal play.

Coding of an octal game (Conway code)

$$d_0 \bullet d_1 d_2 d_3 \cdots \quad d_i \in \{0, \dots, 7\}$$

d_i written in base 8: $e_2^{(i)} e_1^{(i)} e_0^{(i)}$ gives the conditions under which i token may be removed.

- ▶ $e_0^{(i)} = 1$, then a (full) heap with i token can be suppressed
- ▶ $e_1^{(i)} = 1$, then a heap with $n > i$ token can be replaced with a heap with $n - i$ token left
- ▶ $e_2^{(i)} = 1$, then a heap with n token can be replaced with two heaps containing respectively a and b token, $a, b \geq 1$,
 $a + b = n - i$.

EXAMPLE

The game of NIM is coded by $0 \bullet 3333 \cdots$, $\text{rep}_8(3) = 011$.

A finite subtraction game $I = \{3, 5, 6\}$ is of the form $0 \bullet 003033$.

THEOREM (OCTAL GAME PERIODICITY [1])

Consider a finite octal game $d_0 \bullet d_1 d_2 \cdots d_k$. If there exist $N \geq 0$ and $p \geq 1$ such that

$$\mathcal{G}(n + p) = \mathcal{G}(n), \quad \forall N \leq n < 2N + p + \max I$$

then $\mathcal{G}(n + p) = \mathcal{G}(n)$ for all $n \geq N$.

Are all finite octal games ultimately periodic? [55, R. Guy]

$0 \bullet 07$ has period 34 and preperiod 53 [Guy, Smith 1956]

$0 \bullet 007$ **no known periodicity**...

$0 \bullet 165$ has period 1550 and preperiod 5181 [56, Austin, 1976]

$0 \bullet 106$ has period $\simeq 3 \cdot 10^{11}$ and preperiod $\simeq 4 \cdot 10^{11}$
[Flammenkamp, 2002]

<http://wwwhomes.uni-bielefeld.de/achim/octal.html>

With more than one pile (let's say two piles).


NATURAL QUESTION

What is the structure of the \mathcal{G} -values for a finite subtraction game over k piles? Do we get a Presburger definable set, i.e., each value determines a semi-linear set?

Cobham–Semenov' theorem: Let $p, q \geq 2$ be multiplicatively independent integers. If $X \subset \mathbb{N}^k$ is both p -recognizable and q -recognizable, then it is definable by a first-order formula in the Presburger arithmetic $\langle \mathbb{N}, + \rangle$ [10].

Work in progress: X. Badin De Montjoye, V. Gledel, V. Marsault, A. Massuir, M.R.


If we let $I = \{(2, 1), (3, 5)\}$, we get




PROPOSITION

If $\#I = 2$, then the set of \mathcal{G} -values is Presburger definable.


If we let $I = \{(1, 3), (3, 1), (4, 4)\}$, we get



If we let $I = \{(1, 2), (2, 1), (3, 5), (5, 3)\}$, we get




If we let $I = \{(1, 2), (2, 1), (3, 5), (5, 3), (2, 2)\}$, we get




We think that this one is NOT Presburger definable.


If we let $I = \{(10, 2), (2, 10), (32, 5), (5, 32), (10, 10)\}$, we get




If we let $I = \{(10, 2), (2, 10), (32, 5), (5, 32), (10, 10)\}$, we get




Cellular automata — kind of space-time diagram with bounded memory, the rules are (1, 2) and (3, 1)




Cellular automata — kind of space-time diagram with bounded memory, the rules are (1, 2) and (3, 1)




Cellular automata — kind of space-time diagram with bounded memory, the rules are (1, 2) and (3, 1)




Cellular automata — kind of space-time diagram with bounded memory, the rules are (1, 2) and (3, 1)




Cellular automata — kind of space-time diagram with bounded memory, the rules are $(1, 2)$ and $(3, 1)$




Cellular automata — kind of space-time diagram with bounded memory, the rules are (1, 2) and (3, 1)










Cellular automata — kind of space-time diagram with bounded memory, the rules are (1, 2) and (3, 1)















Cellular automata — kind of space-time diagram with bounded memory, the rules are (1, 2) and (3, 1)















Bibliography (in chronological order of the slides)







-  A. Siegel, *Combinatorial game theory*, Graduate Studies in Mathematics, **146**, AMS (2013).
-  E. R. Berlekamp, J. H. Conway, R. K. Guy, *Winning ways for your mathematical plays*, A K Peters, Ltd. (2001).
-  T. S. Ferguson, *Game Theory*, UCLA.
-  C. Berge, *Graphes et hypergraphes*, Monographies Universitaires de Mathématiques, No. 37. Dunod, Paris, (1970).
-  W. A. Wythoff, A modification of the game of Nim, *Nieuw Arch. Wiskd.* **7** (1907), 199–202.
-  C. L. Bouton, Nim, a game with a complete mathematical theory, *Ann. of Math.* (2) **3** (1901/02), no. 1-4, 35–39.
-  J.-P. Allouche, J. Shallit, *Automatic sequences. Theory, applications, generalizations*. Cambridge Univ. Press, Cambridge, (2003).







-  A. Cobham 1972, Uniform tag sequences, *Math. Systems Theory* **6** (1972), 164–192.
-  J.R. Büchi, Weak second-order arithmetic and finite automata, *Z. Math. Logik Grundlag. Math.* **6** (1960), 66–92.
-  V. Bruyère, G. Hansel, C. Michaux, R. Villemaire, Logic and p -recognizable sets of integers, *Bull. Belg. Math. Soc. Simon Stevin* **1** (1994), no. 2, 191–238.
-  M. Rigo, *Formal Languages, Automata and Numeration Systems*, vol. 2, *Applications to recognizability and decidability*. Networks and Telecommunications Series. ISTE, London; John Wiley & Sons, Inc., Hoboken, NJ, (2014).
-  G. Christol, Ensembles presque périodiques k -reconnaissables, *Theoret. Comput. Sci.* **9** (1979), 141–145.
-  G. Christol, T. Kamae, M. Mendès France, G. Rauzy, Suites algébriques, automates et substitutions, *Bull. Soc. Math. France* **108** (1980), 401–419.








-  E. Rowland, R. Yassawi, A characterization of p -automatic sequences as columns of linear cellular automata, *Adv. in Appl. Math.* **63** (2015), 68–89.
-  S. Eilenberg, *Automata, Languages and Machines*, Vol. A, Academic Press, New-York (1974).
-  J.-P. Allouche, J. Shallit, The ring of k -regular sequences, *Theoret. Comput. Sci.* **98** (1991), 163–197.
-  A. Carpi, V. D'Alonzo, On factors of synchronized sequences, *Theoret. Comput. Sci.* **411** (2010), no. 44-46, 3932–3937.
-  J. Berstel, C. Reutenauer, *Noncommutative rational series with applications*, Encyclopedia of Mathematics and its Applications, vol. **137**, Cambridge Univ. Press (2011).
-  A. Thue, Über unendliche Zeichenreihen, *Norske vid. Selsk. Skr. Mat. Nat. Kl.* **7** (1906), 1–22.







-  A. Thue, Über die gegenseitige Lage gleicher Teile gewisser Zeichenreihen, *Norske vid. Selsk. Skr. Mat. Nat. Kl.* **1** (1912), 1–67.
-  M. Lothaire, *Combinatorics on Words*, Encyclopedia of Mathematics and Its Applications, vol. **17**, Addison-Wesley (1983).
-  V. Keränen, Abelian squares are avoidable on 4 letters, ICALP, pp. 41–52 (1992).
-  J. Cassaigne, J. D. Currie, L. Schaeffer, J. Shallit, Avoiding Three Consecutive Blocks of the Same Size and Same Sum, *J. ACM* **61** (2014), no. 2, Art. 10, 17 pp.
-  E. Prouhet, Mémoire sur quelques relations entre les puissances de nombres, *C. R. Acad. Sci. Paris Sér. I* **33** (1851), 225.
-  P. Borwein, C. Ingalls, The Prouhet-Tarry-Escott problem revisited , *Enseign. Math.* **40** (1994), 3–27.

-  J.-P. Allouche, J. Shallit, The ubiquitous Prouhet-Thue-Morse sequence, Sequences and their applications (Singapore, 1998), 1–16, *Springer Ser. Discrete Math. Theor. Comput. Sci.*, Springer, London, (1999).
-  U. Blass, A.S. Fraenkel, The Sprague-Grundy function for Wythoff's game, *Theoret. Comput. Sci.* **75** (1990), no. 3, 311–333.
-  Y. Jiao, On the Sprague-Grundy values of the \mathcal{F} -Wythoff game. *Electron. J. Combin.* **20** (2013).
-  A. Gu, Sprague-Grundy values of the \mathcal{R} -Wythoff game, *Electron. J. Combin.* **22** (2015).
-  M. Weinstein, Invariance of the Sprague-Grundy function for variants of Wythoff's game, *Integers* **16** (2016).
-  A. Dress, A. Flammenkamp and N. Pink, Additive periodicity of the Sprague-Grundy function of certain Nim games, *Adv. in Appl. Math.* **22**, 249–270 (1999).

-  H. A. Landman, A simple FSM⁵-based proof of the additive periodicity of the Sprague-Grundy function of Wythoff's game, *More Games of No Chance*, 2002.
-  S. Beatty, Problem 3173, *American Mathematical Monthly* **33**, 159.
-  M. Lothaire, *Algebraic Combinatorics on Words*, Encyclopedia of Mathematics and its Applications, vol. **90**, Cambridge Univ. Press (2002).
-  É. Zeckendorf, Représentation des nombres naturels par une somme des nombres de Fibonacci ou de nombres de Lucas, *Bull. Soc. Roy. Sci. Liège* **41** (1972) 179–182.
-  J. Shallit, A generalization of automatic sequences, *Theoret. Comput. Sci.* **61** (1988), no. 1, 1–16.
-  J.-P. Allouche, E. Cateland, W. J. Gilbert, H.-O. Peitgen, J. Shallit, G. Skordev, Automatic maps in exotic numeration systems, *Theory Comput. Syst.* **30** (1997), no. 3, 285–331.

-  J.-P. Allouche, K. Scheicher, R. Tichy, Regular maps in generalized number systems, *Math. Slovaca* **50** (2000), no. 1, 41–58.
-  M. Rigo, Generalization of automatic sequences for numeration systems on a regular language, *Theoret. Comput. Sci.* **244** (2000), no. 1-2, 271–281.
-  A. Maes, M. Rigo, More on generalized automatic sequences, *J. Autom. Lang. Comb.* **7** (2002), no. 3, 351–376.
-  Ch. Frougny, On the sequentiality of the successor function, *Inform. and Comput.* **139** (1997) 17–28.
-  A.S. Fraenkel, How to beat your Wythoff games' opponent on three fronts, *Amer. Math. Monthly* **89** (1982), no. 6, 353–361.
-  E. Duchêne, A.S. Fraenkel, R. Nowakowski, M. Rigo, Extensions and restrictions of Wythoff's game preserving its \mathcal{P} -positions, *J. Combin. Theory Ser. A* **117** (2010), no. 5, 545–567.

-  O. Salon, Suites automatiques à multi-indices, *Séminaire de théorie des nombres*, Bordeaux, 1986–1987, exposé 4.
-  A. Maes, Morphisms and almost-periodicity, *Discrete Appl. Math.* **86** (1998), no. 2-3, 233–248.
-  A. Maes, An automata-theoretic decidability proof for first-order theory of $\langle N, <, P \rangle$ with morphic predicate P , *J. Autom. Lang. Comb.* **4** (1999), no. 3, 229–245.
-  A. Maes, *Morphic predicates and applications to the decidability of arithmetic theories*, UMH Univ. Mons-Hainaut, Ph.D. thesis, January 1999.
-  E. Duchêne, M. Rigo, Invariant games, *Theoret. Comput. Sci.* **411** (2010), 3169–3180.
-  E. Duchêne, A. Parreau, M. Rigo, Deciding game invariance, *Inform. and Comput.* **253** (2017), 127–142.
-  P. Lecomte, M. Rigo, Numeration systems on a regular language, *Theory Comput. Syst.* **34** (2001), 27–44.

-  G. Katona, A theorem on finite sets, *Theory of Graphs*, Proc. Colloquium, Tihany, Hungary (1966), 187–207.
-  D. H. Lehmer, The machine tools of combinatorics, in *Applied Combinatorial Mathematics* (E. F. Beckenbach Ed.), Wiley, New York, (1964), 5–31.
-  J. S. Lew, L. B. Morales, A. Sánchez-Flores, Diagonal polynomials for small dimensions, *Math. Systems Theory* **29** (1996), 305–310.
-  É. Charlier, T. Kärki, M. Rigo, Multidimensional generalized automatic sequences and shape-symmetric morphic words, *Discrete Math.* **310** (2010), no. 6-7, 1238–1252.
-  R. K. Guy, C. A. B. Smith, The G -values of various games, *Proc. Cambridge Philos. Soc.* **52** (1956), 514–526.
-  R. B. Austin, *Impartial and partisan games*, Master thesis, Univ. of Calgary (1976).