

# I. Smooth Parametrizations in dynamics

David Burguet

Luminy, May 20, 2019

# Shub's entropy conjecture

$M$  a compact manifold,

$f : M \rightarrow M$  a  $C^0$  map,

$h_{top}$  topological entropy of  $f$ ,

$f_* : H_*(M) \rightarrow H_*(M)$ ,

$\rho$  spectral radius of  $f_*$ .

# Shub's entropy conjecture

$M$  a compact manifold,  
 $f : M \rightarrow M$  a  $\mathcal{C}^0$  map,  
 $h_{top}$  topological entropy of  $f$ ,  
 $f_* : H_*(M) \rightarrow H_*(M)$ ,  
 $\rho$  spectral radius of  $f_*$ .

## Entropy Conjecture (Shub)

$f \in \mathcal{C}^1$ ,

$$h_{top} \geq \log \rho.$$

# Shub's entropy conjecture

$M$  a compact manifold,  
 $f : M \rightarrow M$  a  $\mathcal{C}^0$  map,  
 $h_{top}$  topological entropy of  $f$ ,

$f_* : H_*(M) \rightarrow H_*(M)$ ,  
 $\rho$  spectral radius of  $f_*$ .

## Entropy Conjecture (Shub)

$f \in \mathcal{C}^1$ ,

$$h_{top} \geq \log \rho.$$

## Theorem (Yomdin)

*The conjecture holds true for any  $\mathcal{C}^\infty$  map.*

# Volume growth

$(M, \|\cdot\|)$  compact  $\mathcal{C}^\infty$  Riemannian manifold,  
 $\mathcal{C}^\infty$  disc  $\sigma : (0, 1)^k \rightarrow M$ , i.e. a  $\mathcal{C}^\infty$  map  
with  $\|d^r \sigma\| < +\infty$  for all  $r \in \mathbb{N}$ .

# Volume growth

$(M, \|\cdot\|)$  compact  $\mathcal{C}^\infty$  Riemannian manifold,  
 $\mathcal{C}^\infty$  disc  $\sigma : (0, 1)^k \rightarrow M$ , i.e. a  $\mathcal{C}^\infty$  map  
with  $\|d^r \sigma\| < +\infty$  for all  $r \in \mathbb{N}$ .

Volume growth of  $\sigma$  :

$$\begin{aligned} v(\sigma) &= \limsup_n \frac{1}{n} \log \text{vol}_k(f^n \circ \sigma), \\ &= \limsup_n \frac{1}{n} \log \int_{(0,1)^k} \|\Lambda^k d_t(f^n \circ \sigma)\| dt, \end{aligned}$$

# Volume growth

$(M, \|\cdot\|)$  compact  $C^\infty$  Riemannian manifold,  
 $C^\infty$  disc  $\sigma : (0, 1)^k \rightarrow M$ , i.e. a  $C^\infty$  map  
with  $\|d^r \sigma\| < +\infty$  for all  $r \in \mathbb{N}$ .

Volume growth of  $\sigma$  :

$$\begin{aligned} v(\sigma) &= \limsup_n \frac{1}{n} \log \text{vol}_k(f^n \circ \sigma), \\ &= \limsup_n \frac{1}{n} \log \int_{(0,1)^k} \|\Lambda^k d_t(f^n \circ \sigma)\| dt, \end{aligned}$$

Global volume growth :

$$v = \sup_{\sigma} v(\sigma).$$

# Volume growth

$(M, \|\cdot\|)$  compact  $\mathcal{C}^\infty$  Riemannian manifold,  
 $\mathcal{C}^\infty$  disc  $\sigma : (0, 1)^k \rightarrow M$ , i.e. a  $\mathcal{C}^\infty$  map  
with  $\|d^r \sigma\| < +\infty$  for all  $r \in \mathbb{N}$ .

Volume growth of  $\sigma$  :

$$\begin{aligned} v(\sigma) &= \limsup_n \frac{1}{n} \log \text{vol}_k(f^n \circ \sigma), \\ &= \limsup_n \frac{1}{n} \log \int_{(0,1)^k} \|\Lambda^k d_t(f^n \circ \sigma)\| dt, \end{aligned}$$

Global volume growth :

$$v = \sup_{\sigma} v(\sigma).$$

Fact

$f \in \mathcal{C}^1$ ,

$$\log \rho \leq v.$$



Volume growth of  $\sigma$  at scale  $\epsilon > 0$  :

$$\begin{aligned}v^*(\sigma, \epsilon) &= \limsup_n \frac{1}{n} \log \sup_{x \in M} \text{vol}_k(f^n \circ \sigma|_{\sigma^{-1}B_n(x, \epsilon)}), \\&= \limsup_n \frac{1}{n} \log \sup_{x \in M} \int_{\sigma^{-1}B_n(x, \epsilon)} \|\Lambda^k d_t(f^n \circ \sigma)\| dt,\end{aligned}$$

$$\text{with } B_n(x, \epsilon) := \bigcap_{0 \leq k < n} f^{-k} B(f^k x, \epsilon) \text{ for } x \in M,$$

Volume growth of  $\sigma$  at scale  $\epsilon > 0$  :

$$\begin{aligned} v^*(\sigma, \epsilon) &= \limsup_n \frac{1}{n} \log \sup_{x \in M} \text{vol}_k(f^n \circ \sigma|_{\sigma^{-1}B_n(x, \epsilon)}), \\ &= \limsup_n \frac{1}{n} \log \sup_{x \in M} \int_{\sigma^{-1}B_n(x, \epsilon)} \|\Lambda^k d_t(f^n \circ \sigma)\| dt, \end{aligned}$$

$$\text{with } B_n(x, \epsilon) := \bigcap_{0 \leq l < n} f^{-l} B(f^l x, \epsilon) \text{ for } x \in M,$$

Local volume growth :

$$v^* = \lim_{\epsilon \rightarrow 0} \sup_{\sigma} v^*(\sigma, \epsilon)$$

Fact

$$v \leq h_{top} + v^*.$$

## Fact

$$\nu \leq h_{top} + \nu^*.$$

$\mathcal{M}$  compact set of  $f$ -invariant probas,  
 $h(\nu)$  metric entropy of  $\nu \in \mathcal{M}$

## Theorem (Newhouse)

$f \in \mathcal{C}^{1+}$ ,

- $h_{top} \leq \nu$ ,
- $\forall \mu \in \mathcal{M}, \limsup_{\nu \rightarrow \mu} h(\nu) \leq h(\mu) + \nu^*.$

## Theorem (Yomdin)

$f \in \mathcal{C}^\infty$ ,

$$v^* = 0.$$

## Theorem (Yomdin)

$f \in \mathcal{C}^\infty$ ,

$$v^* = 0.$$

Proof of Entropy Conjecture for  $\mathcal{C}^\infty$  systems :

$$\log \rho \leq v \leq h_{top} + v^* = h_{top}.$$



## Theorem (Yomdin)

$f \in \mathcal{C}^\infty$ ,

$$v^* = 0.$$

Proof of Entropy Conjecture for  $\mathcal{C}^\infty$  systems :

$$\log \rho \leq v \leq h_{top} + v^* = h_{top}.$$



## Corollary

$f \in \mathcal{C}^\infty$ ,

- $h_{top} = v$ ,
- $\forall \mu \in \mathcal{M}$ ,  $\limsup_{\nu \rightarrow \mu} h(\nu) \leq h(\mu)$ . *In particular there exists an equilibrium measure w.r.t. any  $\mathcal{C}^0$  potential.*

$f : M \hookrightarrow \mathbb{R}^d$  a  $C^r$  map with  $+\infty > r \geq 1$ ,  $d = \dim(M)$ ,  
 $\sigma : (0, 1)^k \rightarrow M$  a  $C^r$  disc, i.e. a  $C^r$  map with  $\|d^r \sigma\| < +\infty$ .



$f : M \rightarrow \mathbb{R}$  a  $C^r$  map with  $+\infty > r \geq 1$ ,  $d = \dim(M)$ ,  
 $\sigma : (0, 1)^k \rightarrow M$  a  $C^r$  disc, i.e. a  $C^r$  map with  $\|d^r \sigma\| < +\infty$ .

$$\mu \in \mathcal{M}, \chi_1^+(\mu) = \lim_n \frac{1}{n} \int \log^+ \|d_x f^n\| d\mu(x),$$
$$R(f) = \lim_n \frac{1}{n} \log^+ \|df^n\|,$$

$f : M \rightarrow M$  a  $C^r$  map with  $+\infty > r \geq 1$ ,  $d = \dim(M)$ ,  
 $\sigma : (0, 1)^k \rightarrow M$  a  $C^r$  disc, i.e. a  $C^r$  map with  $\|d^r \sigma\| < +\infty$ .

$$\mu \in \mathcal{M}, \chi_1^+(\mu) = \lim_n \frac{1}{n} \int \log^+ \|d_x f^n\| d\mu(x),$$
$$R(f) = \lim_n \frac{1}{n} \log^+ \|df^n\|,$$


## Theorem

- $\nu^* \leq \frac{kR(f)}{r},$
- $\limsup_{\nu \rightarrow \mu} h(\nu) \leq h(\mu) + \frac{d\chi_1^+(\mu)}{r}.$


These upperbounds are essentially sharp. Moreover there are  $C^r$  examples without maximal measures (Misiurewicz, Buzzi).

$C^r$  example with  $v^* \neq 0$  :  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  given by  $\begin{pmatrix} \lambda^{-1} & 0 \\ 0 & \lambda \end{pmatrix}$  with  $\lambda > 1$

$$\begin{aligned} \sigma : (0, 1) &\rightarrow \mathbb{R}^2, \\ t &\mapsto (t, t^{2r+1} \sin(1/t))', \quad x = 0. \end{aligned}$$




$\sigma \cap [0,1]^2$




$f^n(\sigma \cap B_n(0,1))$

$C^r$  example with  $v^* \neq 0$  :  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  given by  $\begin{pmatrix} \lambda^{-1} & 0 \\ 0 & \lambda \end{pmatrix}$  with  $\lambda > 1$

$$\begin{aligned} \sigma : (0, 1) &\rightarrow \mathbb{R}^2, \\ t &\mapsto (t, t^{2r+1} \sin(1/t)) \quad , \quad x = 0. \end{aligned}$$



$$\sigma \cap [0, 1]^2$$




$$f^n(\sigma \cap B_n(0, 1))$$

$y_{f^n(P)} = x_P^{2r+1} \times \lambda^n \simeq 1$  and  
 $\simeq 1/x_P$  disc. branches in  $f^n(\sigma \cap B_n(0, 1))$


$$v^*(\sigma, 1) \geq \lim_n \frac{\log(1/x_P)}{n} = \frac{\log \lambda}{2r+1}.$$

$\mathcal{C}^r$  example with  $v^* \neq 0$  :  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  given by  $\begin{pmatrix} \lambda^{-1} & 0 \\ 0 & \lambda \end{pmatrix}$  with  $\lambda > 1$

$$\begin{aligned} \sigma : (0, 1) &\rightarrow \mathbb{R}^2, \\ t &\mapsto (t, t^{2r+1} \sin(1/t))', \quad x = 0. \end{aligned}$$



$\sigma \cap [0,1]^2$



$f^n(\sigma \cap B_n(0,1))$

$$v^*(\sigma) \geq \frac{\log \lambda}{2r+1}.$$

$B$  the unit euclidean ball in  $\mathbb{R}^d$ ,

$P : (0, 1)^k \rightarrow \mathbb{R}^d$  with  $P = (P_1, \dots, P_d) \in \mathbb{R}^d[X_1, \dots, X_k]$ ,

$s = \max_i \deg P_i$ ,

$B$  the unit euclidean ball in  $\mathbb{R}^d$ ,

$P : (0, 1)^k \rightarrow \mathbb{R}^d$  with  $P = (P_1, \dots, P_d) \in \mathbb{R}^d[X_1, \dots, X_k]$ ,

$s = \max_i \deg P_i$ ,

$r \in \mathbb{N}$ , for a  $\mathcal{C}^r$  disc  $\varphi : (0, 1)^k \rightarrow \mathbb{R}^d$ ,

we let  $\|\varphi\|_r := \max_{q \leq r} \|d^q \varphi\|$ ,

rep. of  $(0, 1)^k$  is a  $\mathcal{C}^\infty$  map from  $(0, 1)^k$  to itself.

$B$  the unit euclidean ball in  $\mathbb{R}^d$ ,

$P : (0, 1)^k \rightarrow \mathbb{R}^d$  with  $P = (P_1, \dots, P_d) \in \mathbb{R}^d[X_1, \dots, X_k]$ ,

$s = \max_i \deg P_i$ ,

$r \in \mathbb{N}$ , for a  $\mathcal{C}^r$  disc  $\varphi : (0, 1)^k \rightarrow \mathbb{R}^d$ ,

we let  $\|\varphi\|_r := \max_{q \leq r} \|d^q \varphi\|$ ,

rep. of  $(0, 1)^k$  is a  $\mathcal{C}^\infty$  map from  $(0, 1)^k$  to itself.

## Lemma (Gromov)

*There exists a family  $\Theta = \{\theta\}$  of rep. of  $(0, 1)^k$  s.t.*

- ①  $\bigcup_{\theta \in \Theta} \text{Im}(\theta) = P^{-1}(B)$ ,
- ②  $\forall \theta \in \Theta, \|\theta\|_r \leq 1$  and  $\|P \circ \theta\|_r \leq 1$ ,
- ③  $\#\Theta \leq \mathfrak{C}$  with  $\mathfrak{C} = \mathfrak{C}(k, d, r, s)$ .



Lemma (B.-Liao-Yang, Binyamini-Novikov)

*There exists  $R_k \in \mathbb{R}[X, Y]$ , s.t.*

$$\mathfrak{C}(k, d, r, s) = R_{k,d}(r, s).$$

## Lemma (B.-Liao-Yang, Binyamini-Novikov)

*There exists  $R_k \in \mathbb{R}[X, Y]$ , s.t.*

$$\mathfrak{C}(k, d, r, s) = R_{k,d}(r, s).$$

## Theorem (Yomdin, B.-Liao-Yang)

*There is an explicit (essentially sharp) rate of convergence of  $\lim_{\epsilon \rightarrow 0} \sup_{\sigma} v^*(\sigma, \epsilon) = 0$  for  $C^\infty$  maps  $f$  and  $\sigma$  in a given ultradifferentiable class, e.g. in the analytic case*

$$\forall \epsilon > 0, \quad \sup_{\sigma} v^*(\sigma, \epsilon) \leq O(\|df\|) \frac{\log(|\log \epsilon|)}{|\log \epsilon|}.$$

$\mathfrak{s} : (0, 1)^k \rightarrow \mathbb{R}^d$  a  $C^r$  disc with  $r \in \mathbb{N}^*$ .

### Lemma

*There exists a family  $\Theta = \{\theta\}$  of rep. of  $(0, 1)^k$  s.t.*

- $\bigcup_{\theta \in \Theta} \text{Im}(\theta) \supset \mathfrak{s}^{-1}(B),$
- $\forall \theta \in \Theta, \|\theta\|_r \leq 1$  and  $\|\mathfrak{s} \circ \theta\|_r \leq 1,$
- $\#\Theta \leq \mathfrak{D} \max(\|d^r \mathfrak{s}\|, 1)^{\frac{k}{r}}$  with  $\mathfrak{D} = \mathfrak{D}(k, d, r).$

$\mathfrak{s} : (0, 1)^k \rightarrow \mathbb{R}^d$  a  $C^r$  disc with  $r \in \mathbb{N}^*$ .

### Lemma

*There exists a family  $\Theta = \{\theta\}$  of rep. of  $(0, 1)^k$  s.t.*

- $\bigcup_{\theta \in \Theta} \text{Im}(\theta) \supset \mathfrak{s}^{-1}(B),$
- $\forall \theta \in \Theta, \|\theta\|_r \leq 1$  and  $\|\mathfrak{s} \circ \theta\|_r \leq 1,$
- $\#\Theta \leq \mathfrak{D} \max(\|d^r \mathfrak{s}\|, 1)^{\frac{k}{r}}$  with  $\mathfrak{D} = \mathfrak{D}(k, d, r).$

Proof : We may assume  $\|d^r \mathfrak{s}\| \leq 1$  : consider  $k$ -subcubes  $C$  of  $(0, 1)^k$  of size  $|C| = \max(\|d^r \mathfrak{s}\|, 1)^{-1/r}$  covering  $(0, 1)^k$  and  $\psi_C : (0, 1)^k \rightarrow C$  affine parametrization of  $C$ , then  $\|d^r(\mathfrak{s} \circ \psi_C)\| = |C|^r \|d^r \mathfrak{s}\| \leq 1 \dots$

$\mathfrak{s} : (0, 1)^k \rightarrow \mathbb{R}^d$  a  $\mathcal{C}^r$  disc with  $r \in \mathbb{N}^*$ .

### Lemma

There exists a family  $\Theta = \{\theta\}$  of rep. of  $(0, 1)^k$  s.t.

- $\bigcup_{\theta \in \Theta} \text{Im}(\theta) \supset \mathfrak{s}^{-1}(B)$ ,
- $\forall \theta \in \Theta, \|\theta\|_r \leq 1$  and  $\|\mathfrak{s} \circ \theta\|_r \leq 1$ ,
- $\#\Theta \leq \mathfrak{D} \max(\|d^r \mathfrak{s}\|, 1)^{\frac{k}{r}}$  with  $\mathfrak{D} = \mathfrak{D}(k, d, r)$ .

Proof : We may assume  $\|d^r \mathfrak{s}\| \leq 1$  : consider  $k$ -subcubes  $C$  of  $(0, 1)^k$  of size  $|C| = \max(\|d^r \mathfrak{s}\|, 1)^{-1/r}$  covering  $(0, 1)^k$  and  $\psi_C : (0, 1)^k \rightarrow C$  affine parametrization of  $C$ , then  $\|d^r(\mathfrak{s} \circ \psi_C)\| = |C|^r \|d^r \mathfrak{s}\| \leq 1 \dots$

If  $\|d^r \mathfrak{s}\| \leq 1$ , let  $P$  be the  $(r-1)$ -Lagrange polynomial of  $\mathfrak{s}$  at  $x_0 \in (0, 1)^k$  and  $\Theta = \{\theta\}$  as in the Algebraic RL for  $\frac{P}{2}$ , then

- $\mathfrak{s}^{-1}(B) \subset P^{-1}(2B) = \bigcup_{\theta \in \Theta} \text{Im}(\theta)$ ,
- $\|\mathfrak{s} \circ \theta\|_r \leq \|P \circ \theta\|_r + \|(\mathfrak{s} - P) \circ \theta\|_r \leq \mathfrak{E} = \mathfrak{E}(k, d, r)$ .

# DRL for non autonomous $C^r$ dynamical systems

$\mathcal{F} = (f_m)_{m \in \mathbb{N}^*}$  family of  $C^r$  maps from  $B$  to  $\mathbb{R}^d$  with  $f_0 = \text{Id}_B$ ,  
 $f^m = f_m \circ \cdots \circ f_0$  from  $B_m$  to  $\mathbb{R}^d$  with  
 $B_m := \bigcap_{0 \leq l < m} f^{-l} B$  dynamical ball,  
 $\mathfrak{s} : (0, 1)^k \rightarrow \mathbb{R}^d$  a  $C^r$  disc.

# DRL for non autonomous $C^r$ dynamical systems

$\mathcal{F} = (f_m)_{m \in \mathbb{N}^*}$  family of  $C^r$  maps from  $B$  to  $\mathbb{R}^d$  with  $f_0 = \text{Id}_B$ ,  
 $f^m = f_m \circ \dots \circ f_0$  from  $B_m$  to  $\mathbb{R}^d$  with  
 $B_m := \bigcap_{0 \leq l < m} f^{-l} B$  dynamical ball,  
 $\mathfrak{s} : (0, 1)^k \rightarrow \mathbb{R}^d$  a  $C^r$  disc.

## Lemma

For any  $m \in \mathbb{N}^*$  there exists  $\Theta_m = \{\theta_m\}$  family of rep. of  $(0, 1)^k$  satisfying :

- $\bigcup_{\theta_m \in \Theta_m} \text{Im}(\theta_m) \supset \mathfrak{s}^{-1}(B_m),$
- $\forall \theta_m \in \Theta_m \forall 0 \leq l < m, \|f^l \circ \mathfrak{s} \circ \theta_m\|_r \leq 1,$
- $\#\Theta_m \leq \mathfrak{D}^m \max(\|d^r \mathfrak{s}\|, 1)^{\frac{k}{r}} \prod_{l=0}^{m-1} \max(\|f_l\|_r, 1)^{\frac{k}{r}}.$

# DRL for non autonomous $C^r$ dynamical systems

$\mathcal{F} = (f_m)_{m \in \mathbb{N}^*}$  family of  $C^r$  maps from  $B$  to  $\mathbb{R}^d$  with  $f_0 = \text{Id}_B$ ,  
 $f^m = f_m \circ \dots \circ f_0$  from  $B_m$  to  $\mathbb{R}^d$  with  
 $B_m := \bigcap_{0 \leq l < m} f^{-l} B$  dynamical ball,  
 $\mathfrak{s} : (0, 1)^k \rightarrow \mathbb{R}^d$  a  $C^r$  disc.

## Lemma

For any  $m \in \mathbb{N}^*$  there exists  $\Theta_m = \{\theta_m\}$  family of rep. of  $(0, 1)^k$  satisfying :

- $\bigcup_{\theta_m \in \Theta_m} \text{Im}(\theta_m) \supset \mathfrak{s}^{-1}(B_m)$ ,
- $\forall \theta_m \in \Theta_m \forall 0 \leq l < m, \|f^l \circ \mathfrak{s} \circ \theta_m\|_r \leq 1$ ,
- $\#\Theta_m \leq \mathfrak{D}^m \max(\|d^r \mathfrak{s}\|, 1)^{\frac{k}{r}} \prod_{l=0}^{m-1} \max(\|f_l\|_r, 1)^{\frac{k}{r}}$ .

Proof : By induction. For each  $\theta_m \in \Theta_m$ , let  $\Theta(\theta_m)$  be the family of rep. obtained when applying the above Lemma to  $f^m \circ \mathfrak{s} \circ \theta_m$ .  
Take  $\Theta_{m+1} = \{\theta_m \circ \theta \mid \theta_m \in \Theta_m, \theta \in \Theta(\theta_m)\}$ .



$\Phi = (\phi_m)_{m \in \mathbb{N}}$  family of  $\alpha$ -Hölder maps from  $B$  to  $\mathbb{R}$   
with  $\sup_m |\phi_m|_\alpha \leq 1$  for some  $0 < \alpha \leq 1$ ,  
 $S_m \Phi = \sum_{l=0}^{m-1} \phi_l \circ f^l$ .

$\Phi = (\phi_m)_{m \in \mathbb{N}}$  family of  $\alpha$ -Hölder maps from  $B$  to  $\mathbb{R}$   
 with  $\sup_m |\phi_m|_\alpha \leq 1$  for some  $0 < \alpha \leq 1$ ,  
 $S_m \Phi = \sum_{l=0}^{m-1} \phi_l \circ f^l$ .

### Lemma

For any  $m \in \mathbb{N}^*$  there exists  $\Theta_m = \{\theta_m\}$  family of rep. of  $(0, 1)^k$  satisfying :

- $\bigcup_{\theta_m \in \Theta_m} \text{Im}(\theta_m) \supset \mathfrak{s}^{-1}(B_m)$ ,
- $\forall \theta_m \in \Theta_m \forall 0 \leq l < m, \|f^l \circ \mathfrak{s} \circ \theta_m\|_r \leq 1$ ,
- $\forall \theta_m \in \Theta_m, |S_m \Phi \circ \mathfrak{s} \circ \theta_m|_\alpha \leq 1$ ,
- $\#\Theta_m \leq m^{1/\alpha} \mathfrak{D}^m \max(\|d^r \mathfrak{s}\|, 1)^{\frac{k}{r}} \prod_{l=0}^{m-1} \max(\|f_l\|_r, 1)^{\frac{k}{r}}$ .

$\Phi = (\phi_m)_{m \in \mathbb{N}}$  family of  $\alpha$ -Hölder maps from  $B$  to  $\mathbb{R}$   
 with  $\sup_m |\phi_m|_\alpha \leq 1$  for some  $0 < \alpha \leq 1$ ,  
 $S_m \Phi = \sum_{l=0}^{m-1} \phi_l \circ f^l$ .

### Lemma

For any  $m \in \mathbb{N}^*$  there exists  $\Theta_m = \{\theta_m\}$  family of rep. of  $(0, 1)^k$  satisfying :

- $\bigcup_{\theta_m \in \Theta_m} \text{Im}(\theta_m) \supset \mathfrak{s}^{-1}(B_m)$ ,
- $\forall \theta_m \in \Theta_m \forall 0 \leq l < m, \|f^l \circ \mathfrak{s} \circ \theta_m\|_r \leq 1$ ,
- $\forall \theta_m \in \Theta_m, |S_m \Phi \circ \mathfrak{s} \circ \theta_m|_\alpha \leq 1$ ,
- $\#\Theta_m \leq m^{1/\alpha} \mathfrak{D}^m \max(\|d^r \mathfrak{s}\|, 1)^{\frac{k}{r}} \prod_{l=0}^{m-1} \max(\|f_l\|_r, 1)^{\frac{k}{r}}$ .

Proof : Let  $\theta'_m$  be the rep. of the previous Lemma, then

$|\phi_l \circ f^l \circ \mathfrak{s} \circ \theta'_m|_\alpha \leq |\phi_l|_\alpha \|f^l \circ \mathfrak{s} \circ \theta'_m\|_1^\alpha \leq 1$ , thus

$|S_m \Phi \circ \mathfrak{s} \circ \theta'_m|_\alpha \leq m$ . Take finally  $\theta_m = \theta'_m \circ \psi_C$  with  $|C| = m^{-1/\alpha}$ .

# DRL for $\mathcal{C}^\infty$ systems

$f : M \rightarrow \mathbb{R}$  a  $\mathcal{C}^\infty$  map,

$\sigma : (0, 1)^k \rightarrow M$  a  $\mathcal{C}^\infty$  disc,

$x \in M$ .

# DRL for $\mathcal{C}^\infty$ systems

$f : M \rightarrow M$  a  $\mathcal{C}^\infty$  map,  
 $\sigma : (0, 1)^k \rightarrow M$  a  $\mathcal{C}^\infty$  disc,  
 $x \in M$ .

$\forall \gamma > 0, \exists \epsilon = \epsilon(f, \gamma)$  and  $C = C(f, \sigma, \gamma) > 0$  s.t.

## Lemma

*For any  $n \in \mathbb{N}^*$  there exists  $\Theta_n = \{\theta_n\}$  family of rep. of  $(0, 1)^k$  satisfying :*

- $\bigcup_{\theta_n \in \Theta_n} \text{Im}(\theta_n) \supset \sigma^{-1}B_n(x, \epsilon),$
- $\forall \theta_n \in \Theta_n \forall 0 \leq l < n, \|d(f^l \circ \sigma \circ \theta_n)\| \leq 1,$
- $\#\Theta_n \leq Ce^{\gamma n}.$

$f : M \rightarrow M$  a  $\mathcal{C}^\infty$  map,  
 $\sigma : (0, 1)^k \rightarrow M$  a  $\mathcal{C}^\infty$  disc,  
 $x \in M$ .

$\forall \gamma > 0, \exists \epsilon = \epsilon(f, \gamma)$  and  $C = C(f, \sigma, \gamma) > 0$  s.t.

## Lemma

For any  $n \in \mathbb{N}^*$  there exists  $\Theta_n = \{\theta_n\}$  family of rep. of  $(0, 1)^k$  satisfying :

- $\bigcup_{\theta_n \in \Theta_n} \text{Im}(\theta_n) \supset \sigma^{-1}B_n(x, \epsilon),$
- $\forall \theta_n \in \Theta_n \forall 0 \leq l < n, \|d(f^l \circ \sigma \circ \theta_n)\| \leq 1,$
- $\#\Theta_n \leq Ce^{\gamma n}.$

Proof of Yomdin's theorem :

$$\sup_{\sigma} v^*(\sigma, \epsilon) \leq \gamma \text{ and then } v^* = 0.$$

$\phi : M \rightarrow \mathbb{R}$  a  $\alpha$ -Hölder potential with  $0 < \alpha \leq 1$ ,

$$S_n \phi = \sum_{l=0}^{n-1} \phi \circ f^l,$$

$\forall \gamma > 0, \exists \epsilon = \epsilon(f, \phi, \gamma)$  and  $C = C(f, \phi, \sigma, \gamma) > 0$  s.t.

### Lemma

*For any  $n \in \mathbb{N}^*$  there exists  $\Theta_n = \{\theta_n\}$  family of rep. of  $(0, 1)^k$  satisfying :*

- $\bigcup_{\theta_n \in \Theta_n} \text{Im}(\theta_n) \supset \sigma^{-1} B_n(x, \epsilon),$
- $\forall \theta_n \in \Theta_n \forall 0 \leq l < n, \|d(f^l \circ \sigma \circ \theta_n)\| \leq 1,$
- $\forall \theta_n \in \Theta_n, \forall t, s \in \text{Im}(\theta_n), |S_n \phi \circ \sigma(t) - S_n \phi \circ \sigma(s)| \leq 1,$
- $\#\Theta_n \leq C e^{\gamma n}.$

Remark : Third item may be seen as a weak Bowen property for  $\phi$  :

$$\exists \epsilon > 0 \exists C > 0 \text{ s.t.}$$

$$\forall n \in \mathbb{N} \forall y \in B_n(x, \epsilon), |S_n \phi(x) - S_n \phi(y)| < C.$$

# Local dynamics of a $C^\infty$ system $(f, M)$ with $\phi : M \rightarrow \mathbb{R}$ a $\alpha$ -Hölder potential

$M = \mathbb{R}^d / \mathbb{Z}^d$ ,  $x \in \mathbb{R}^d$  and  $\bar{x} \in \mathbb{R}^d / \mathbb{Z}^d$  fixed,  
 $\psi_{\bar{x}}^\epsilon = \overline{x + \epsilon \cdot}$  from  $\mathbb{R}^d$  to  $\mathbb{R}^d / \mathbb{Z}^d$ ,



# Local dynamics of a $C^\infty$ system $(f, M)$ with $\phi : M \rightarrow \mathbb{R}$ a $\alpha$ -Hölder potential

$M = \mathbb{R}^d / \mathbb{Z}^d$ ,  $x \in \mathbb{R}^d$  and  $\bar{x} \in \mathbb{R}^d / \mathbb{Z}^d$  fixed,

$\psi_{\bar{x}}^\epsilon = \overline{x + \epsilon \cdot}$  from  $\mathbb{R}^d$  to  $\mathbb{R}^d / \mathbb{Z}^d$ ,

$\sigma : (0, 1)^k \rightarrow M$  a  $C^\infty$  disc with  $\text{Im}(\sigma) \subset B(x, 2\epsilon)$ ,

$\mathfrak{s} = (\psi_{\bar{x}}^\epsilon)^{-1} \circ \sigma : (0, 1)^k \rightarrow \mathbb{R}^d$ ,

# Local dynamics of a $C^\infty$ system $(f, M)$ with $\phi : M \rightarrow \mathbb{R}$ a $\alpha$ -Hölder potential

$M = \mathbb{R}^d / \mathbb{Z}^d$ ,  $x \in \mathbb{R}^d$  and  $\bar{x} \in \mathbb{R}^d / \mathbb{Z}^d$  fixed,

$\psi_{\bar{x}}^\epsilon = \overline{x + \epsilon \cdot}$  from  $\mathbb{R}^d$  to  $\mathbb{R}^d / \mathbb{Z}^d$ ,

$\sigma : (0, 1)^k \rightarrow M$  a  $C^\infty$  disc with  $\text{Im}(\sigma) \subset B(x, 2\epsilon)$ ,

$\mathfrak{s} = (\psi_{\bar{x}}^\epsilon)^{-1} \circ \sigma : (0, 1)^k \rightarrow \mathbb{R}^d$ ,

$\mathcal{F} = (f_m)_m$  with  $f_m = \left( \psi_{f^p(m+1)\bar{x}}^\epsilon \right)^{-1} \circ f^p \circ \psi_{f^p m \bar{x}}^\epsilon$ ,

$\Phi = (\phi_m)_m$  with  $\phi_m = \sum_{k=0}^{p-1} \phi \circ f^k \circ \psi_{f^p m \bar{x}}^\epsilon$ ,

# Local dynamics of a $C^\infty$ system $(f, M)$ with $\phi : M \rightarrow \mathbb{R}$ a $\alpha$ -Hölder potential

$M = \mathbb{R}^d / \mathbb{Z}^d$ ,  $x \in \mathbb{R}^d$  and  $\bar{x} \in \mathbb{R}^d / \mathbb{Z}^d$  fixed,

$\psi_{\bar{x}}^\epsilon = \overline{x + \epsilon \cdot}$  from  $\mathbb{R}^d$  to  $\mathbb{R}^d / \mathbb{Z}^d$ ,

$\sigma : (0, 1)^k \rightarrow M$  a  $C^\infty$  disc with  $\text{Im}(\sigma) \subset B(x, 2\epsilon)$ ,

$\mathfrak{s} = (\psi_{\bar{x}}^\epsilon)^{-1} \circ \sigma : (0, 1)^k \rightarrow \mathbb{R}^d$ ,

$\mathcal{F} = (f_m)_m$  with  $f_m = \left( \psi_{f^p(m+1)\bar{x}}^\epsilon \right)^{-1} \circ f^p \circ \psi_{f^p m \bar{x}}^\epsilon$ ,

$\Phi = (\phi_m)_m$  with  $\phi_m = \sum_{k=0}^{p-1} \phi \circ f^k \circ \psi_{f^p m \bar{x}}^\epsilon$ ,

$$\forall s \geq 1, \sup_m \|d^s f_m\| = O(\epsilon^{s-1})$$

$$\text{and } \sup_m |\phi_m|_\alpha = O(\epsilon^\alpha) \text{ uniformly in } x.$$

Proof of DRL for  $f : M \curvearrowright \mathcal{C}^\infty$  and  $\phi : M \rightarrow \mathbb{R}$   $\alpha$ -Hölder :

Choose  $r, p, \epsilon$  w.r.t. small error term  $\gamma > 0$  s.t.

- $r \in \mathbb{N}$  with  $\|df\|^{k/r} < e^{\gamma/2}$ ,
- $p \in \mathbb{N}$  with  $\mathfrak{D}^{1/p} < e^{\gamma/2}$ ,
- $\epsilon > 0$  with  $2\epsilon \max(\|df^p\|, 1) < 1$ ,  $\|f_m\|_r \leq \|df^p\|$  and  $|\phi_m|_\alpha \leq 1$  for all  $m \in \mathbb{N}$  with  $\mathcal{F} = (f_m)_m$  and  $(\phi_m)_m$  as above.

Proof of DRL for  $f : M \curvearrowright \mathcal{C}^\infty$  and  $\phi : M \rightarrow \mathbb{R}$   $\alpha$ -Hölder :

Choose  $r, p, \epsilon$  w.r.t. small error term  $\gamma > 0$  s.t.

- $r \in \mathbb{N}$  with  $\|df\|^{k/r} < e^{\gamma/2}$ ,
- $p \in \mathbb{N}$  with  $\mathfrak{D}^{1/p} < e^{\gamma/2}$ ,
- $\epsilon > 0$  with  $2\epsilon \max(\|df^p\|, 1) < 1$ ,  $\|f_m\|_r \leq \|df^p\|$  and  $|\phi_m|_\alpha \leq 1$  for all  $m \in \mathbb{N}$  with  $\mathcal{F} = (f_m)_m$  and  $(\phi_m)_m$  as above.

For  $n = pm$ , we have

$$B_n^f(\bar{x}, \epsilon) \subset \psi_{\bar{x}}^\epsilon(B_m(\mathcal{F})) \text{ and}$$

$$S_n \phi = S_m \Phi \circ (\psi_{\bar{x}}^\epsilon)^{-1}.$$

Proof of DRL for  $f : M \curvearrowright \mathcal{C}^\infty$  and  $\phi : M \rightarrow \mathbb{R}$   $\alpha$ -Hölder :

Choose  $r, p, \epsilon$  w.r.t. small error term  $\gamma > 0$  s.t.

- $r \in \mathbb{N}$  with  $\|df\|^{k/r} < e^{\gamma/2}$ ,
- $p \in \mathbb{N}$  with  $\mathfrak{D}^{1/p} < e^{\gamma/2}$ ,
- $\epsilon > 0$  with  $2\epsilon \max(\|df^p\|, 1) < 1$ ,  $\|f_m\|_r \leq \|df^p\|$  and  $|\phi_m|_\alpha \leq 1$  for all  $m \in \mathbb{N}$  with  $\mathcal{F} = (f_m)_m$  and  $(\phi_m)_m$  as above.

For  $n = pm$ , we have

$$B_n^f(\bar{x}, \epsilon) \subset \psi_{\bar{x}}^\epsilon(B_m(\mathcal{F})) \text{ and}$$

$$S_n \phi = S_m \Phi \circ (\psi_{\bar{x}}^\epsilon)^{-1}.$$

Let  $(\Theta_m(\mathcal{F}))_m$  be the families of rep. given by DRL for n.a.  $\mathcal{C}^r$  systems applied to  $\mathcal{F}$ . The family  $\Theta'_n = \{\theta'_n = \theta_m, \theta_m \in \Theta_m(\mathcal{F})\}$  satisfies DRL for  $f$  :

$$\begin{aligned} \sharp \Theta'_n &\leq m^{1/\alpha} \mathfrak{D}^m \prod_{l=0}^{m-1} \max(\|f_l\|_r, 1)^{\frac{k}{r}}, \\ &\leq m^{1/\alpha} \mathfrak{D}^m \max(\|df\|, 1)^{n \frac{k}{r}}, \\ &\leq Ce^{\gamma n}. \end{aligned}$$

# Asymptotic $h$ -expansiveness of $\mathcal{C}^\infty$ systems

$(X, T)$  top. system, i.e.  $(X, d)$  compact metric space  
and  $T : X \rightarrow X$  continuous,

$n \in \mathbb{N}$ ,  $\delta > 0$ ,  $K \subset X$ ,

$$r_n(\delta, K) = \min \left\{ \#E_\delta, \bigcup_{x \in E_\delta} B_n(x, \delta) \supset K \right\}.$$

Tail entropy of  $T$  :

$$h^* = \lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \limsup_n \frac{1}{n} \log \sup_{x \in X} r_n(\delta, B_n(x, \epsilon)).$$

# Asymptotic $h$ -expansiveness of $\mathcal{C}^\infty$ systems

$(X, T)$  top. system, i.e.  $(X, d)$  compact metric space  
and  $T : X \rightarrow X$  continuous,

$n \in \mathbb{N}$ ,  $\delta > 0$ ,  $K \subset X$ ,

$$r_n(\delta, K) = \min \left\{ \#E_\delta, \bigcup_{x \in E_\delta} B_n(x, \delta) \supset K \right\}.$$

Tail entropy of  $T$  :

$$h^* = \lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \limsup_n \frac{1}{n} \log \sup_{x \in X} r_n(\delta, B_n(x, \epsilon)).$$

## Theorem (Misiurewicz)

$$\forall \mu \in \mathcal{M}, \limsup_{\nu \rightarrow \mu} h(\nu) \leq h(\mu) + h^*.$$



$(X, T)$  is said to be *asymptotically  $h$ -expansive* when  $h^* = 0$ .

$(X, T)$  is said to be *asymptotically  $h$ -expansive* when  $h^* = 0$ .

### Theorem (Buzzi)

*Any  $C^\infty$  system is asymptotically  $h$ -expansive.*

$(X, T)$  is said to be *asymptotically  $h$ -expansive* when  $h^* = 0$ .

### Theorem (Buzzi)

*Any  $C^\infty$  system is asymptotically  $h$ -expansive.*

Proof : With the notations of DRL, take  $\sigma = \psi_X^\epsilon$ . If  $F_\delta$  is  $\delta$  dense in  $(0, 1)^k$  then  $E_\delta = \bigcup_{\theta_n \in \Theta_n} \theta_n(F_\delta)$  is  $\delta$ -dense for the distance  $d_n$  in  $B_n(x, \epsilon)$  with  $\forall x, y \in M$ ,  $d_n(x, y) = \max_{0 \leq k < n} d(f^k x, f^k y)$ .

$(X, T)$  is said to be *asymptotically  $h$ -expansive* when  $h^* = 0$ .

### Theorem (Buzzi)

*Any  $C^\infty$  system is asymptotically  $h$ -expansive.*

Proof : With the notations of DRL, take  $\sigma = \psi_X^\epsilon$ . If  $F_\delta$  is  $\delta$  dense in  $(0, 1)^k$  then  $E_\delta = \bigcup_{\theta_n \in \Theta_n} \theta_n(F_\delta)$  is  $\delta$ -dense for the distance  $d_n$  in  $B_n(x, \epsilon)$  with  $\forall x, y \in M$ ,  $d_n(x, y) = \max_{0 \leq k < n} d(f^k x, f^k y)$ .  
Therefore

$$\begin{aligned} \forall \delta > 0, \quad r_n(\delta, B_n(x, \epsilon)) &\leq \#\Theta_n \times \#F_\delta, \\ &\leq Ce^{\gamma n} \#F_\delta \end{aligned}$$

and

$$h^* \leq \gamma.$$

# DRL for $\mathcal{C}^\infty$ Cocycles

$f : M \rightarrow \mathbb{R}$  a  $\mathcal{C}^\infty$  map,

$\pi : V \rightarrow M$  a  $\mathcal{C}^\infty$  Riemannian vector bundle over  $M$ ,

$F : V \rightarrow V$  a  $\mathcal{C}^\infty$  *semi-invertible* bundle morphism with  $\pi \circ F = f \circ \pi$ ,

$SF : S(V) \rightarrow S(V)$  associated sphere bundle morphism,

$\phi : S(V) \rightarrow \mathbb{R}$  a  $\alpha$ -Hölder potential with  $0 < \alpha \leq 1$ ,

$\Gamma : (0, 1)^k \rightarrow S(V)$  a  $\mathcal{C}^\infty$  disc,  $\sigma = \pi \circ \Gamma$ ,

$x \in M$ .

# DRL for $\mathcal{C}^\infty$ Cocycles

$f : M \curvearrowright$  a  $\mathcal{C}^\infty$  map,

$\pi : V \rightarrow M$  a  $\mathcal{C}^\infty$  Riemannian vector bundle over  $M$ ,

$F : V \curvearrowright$  a  $\mathcal{C}^\infty$  *semi-invertible* bundle morphism with  $\pi \circ F = f \circ \pi$ ,

$SF : S(V) \curvearrowright$  associated sphere bundle morphism,

$\phi : S(V) \rightarrow \mathbb{R}$  a  $\alpha$ -Hölder potential with  $0 < \alpha \leq 1$ ,

$\Gamma : (0, 1)^k \rightarrow S(V)$  a  $\mathcal{C}^\infty$  disc,  $\sigma = \pi \circ \Gamma$ ,

$x \in M$ .

$\forall \gamma > 0, \exists \epsilon = \epsilon(F, \phi, \gamma)$  and  $C = C(F, \phi, \Gamma, \gamma) > 0$  s.t.

## Lemma

For any  $n \in \mathbb{N}^*$  there exists  $\Theta_n = \{\theta_n\}$  family of rep. of  $(0, 1)^k$  satisfying :

- $\bigcup_{\theta_n \in \Theta_n} \text{Im}(\theta_n) \supset \sigma^{-1} B_n^f(x, \epsilon),$
- $\forall \theta_n \in \Theta_n \forall 0 \leq l < n, \|d(SF^l \circ \Gamma \circ \theta_n)\| \leq 1,$
- $\forall \theta_n \in \Theta_n \forall t, s \in \text{Im}(\theta_n), |S_n \phi(\Gamma(t)) - S_n \phi(\Gamma(s))| \leq 1,$
- $\#\Theta_n \leq C e^{\gamma n}.$

## Lemma

For  $\gamma > 0$  and  $x \in M$  there exist  $\tilde{\epsilon} = \tilde{\epsilon}(\gamma)$  and  $C = C(\gamma)$  constant s.t. we have for all  $n \in \mathbb{N}$  and for all  $0 < \epsilon < \tilde{\epsilon}$  :

$$r_n^{\mathbb{S}(F)} \left( \epsilon, \pi^{-1} B_n^f(x, \epsilon) \right) < C e^{\gamma n}.$$

## Lemma

For  $\gamma > 0$  and  $x \in M$  there exist  $\tilde{\epsilon} = \tilde{\epsilon}(\gamma)$  and  $C = C(\gamma)$  constant s.t. we have for all  $n \in \mathbb{N}$  and for all  $0 < \epsilon < \tilde{\epsilon}$  :

$$r_n^{\mathbb{S}(F)} \left( \epsilon, \pi^{-1} B_n^f(x, \epsilon) \right) < C e^{\gamma n}.$$

Proof :  $\gamma > 0$  fixed,

- Sacksteder-Shub : Let  $\mathcal{F} = \{\mathbb{S}(A_i)\}_{i \in \mathbb{N}}$  with  $A_i \in GL(\mathbb{R}^{e+1})$ ,  
 $\forall \epsilon' > 0 \forall n \in \mathbb{N}, r_n^{\mathcal{F}}(\epsilon', \mathbb{S}^e) \leq C_{\epsilon'} n^e$ ,



## Lemma

For  $\gamma > 0$  and  $x \in M$  there exist  $\tilde{\epsilon} = \tilde{\epsilon}(\gamma)$  and  $C = C(\gamma)$  constant s.t. we have for all  $n \in \mathbb{N}$  and for all  $0 < \epsilon < \tilde{\epsilon}$  :

$$r_n^{\mathbb{S}(F)} \left( \epsilon, \pi^{-1} B_n^f(x, \epsilon) \right) < C e^{\gamma n}.$$

Proof :  $\gamma > 0$  fixed,

- Sacksteder-Shub : Let  $\mathcal{F} = \{\mathbb{S}(A_i)\}_{i \in \mathbb{N}}$  with  $A_i \in GL(\mathbb{R}^{e+1})$ ,  
 $\forall \epsilon' > 0 \forall n \in \mathbb{N}, r_n^{\mathcal{F}}(\epsilon', \mathbb{S}^e) \leq C_{\epsilon'} n^e$ ,
- Bowen :  $\forall \epsilon' > 0 \exists \tilde{\epsilon} = \tilde{\epsilon}(\gamma, \epsilon') > 0 \exists C = C(\gamma, \epsilon') > 0$  s.t.  
 $\forall \epsilon < \tilde{\epsilon} \forall n \in \mathbb{N}, \sup_{x \in M} r_n^{\mathbb{S}(F)}(\epsilon', \pi^{-1} B_n^f(x, \epsilon)) \leq C e^{\gamma n/2}$ ,

## Lemma

For  $\gamma > 0$  and  $x \in M$  there exist  $\tilde{\epsilon} = \tilde{\epsilon}(\gamma)$  and  $C = C(\gamma)$  constant s.t. we have for all  $n \in \mathbb{N}$  and for all  $0 < \epsilon < \tilde{\epsilon}$  :

$$r_n^{\mathbb{S}(F)} \left( \epsilon, \pi^{-1} B_n^f(x, \epsilon) \right) < C e^{\gamma n}.$$

Proof :  $\gamma > 0$  fixed,

- Sacksteder-Shub : Let  $\mathcal{F} = \{\mathbb{S}(A_i)\}_{i \in \mathbb{N}}$  with  $A_i \in GL(\mathbb{R}^{e+1})$ ,  
 $\forall \epsilon' > 0 \forall n \in \mathbb{N}, r_n^{\mathcal{F}}(\epsilon', \mathbb{S}^e) \leq C_{\epsilon'} n^e$ ,
- Bowen :  $\forall \epsilon' > 0 \exists \tilde{\epsilon} = \tilde{\epsilon}(\gamma, \epsilon') > 0 \exists C = C(\gamma, \epsilon') > 0$  s.t.  
 $\forall \epsilon < \tilde{\epsilon} \forall n \in \mathbb{N}, \sup_{x \in M} r_n^{\mathbb{S}(F)}(\epsilon', \pi^{-1} B_n^f(x, \epsilon)) \leq C e^{\gamma n/2}$ ,
- Buzzi : Take  $\epsilon' = \epsilon'(\gamma) > 0$  s.t.  
 $\forall \delta > 0, \limsup_n \frac{1}{n} \log \sup_{y \in \mathbb{S}(V)} r_n^{\mathbb{S}(F)}(\delta, B_n^{\mathbb{S}(F)}(y, \epsilon')) < \gamma/2$ .