

Partial Regularity in Time for the Landau Equation (with Coulomb Interaction)

François Golse

CMLS, École polytechnique, Paris

CIRM, March 22-26 2021

Kinetic Equations: from Modeling, Computation to Analysis

Work in collaboration with M.P. Gualdani, C. Imbert and A. Vasseur
to appear in Ann. Sci. Éc. Norm. Sup.
[arXiv:1906.02841 \[math.AP\]](https://arxiv.org/abs/1906.02841)

Landau Equation

Landau equation with unknown $f \equiv f(t, v) \geq 0$:

$$\partial_t f(t, v) = \operatorname{div}_v \int_{\mathbf{R}^3} a(v-w)(\nabla_v - \nabla_w)(f(t, v)f(t, w))dw, \quad v \in \mathbf{R}^3$$

with the notation:

$$a(z) := \frac{1}{8\pi} \nabla^2 |z| = \frac{1}{8\pi|z|} \Pi(z), \quad \Pi(z) := I - \left(\frac{z}{|z|}\right)^{\otimes 2}$$

Nonconservative form

$$\partial_t f(t, v) = (a_{ij} \star_v f(t, v)) \partial_{v_i} \partial_{v_j} f(t, v) + f(t, v)^2$$

Open question global existence of classical solutions or finite-time blow-up for the Cauchy problem with $f|_{t=0} = f_{in}$?

Villani's H-Solutions

H-solution $0 \leq f \in C([0, T); \mathcal{D}'(\mathbf{R}^3)) \cap L^1((0, T); L_{-1}^1(\mathbf{R}^3))$ s.t.

$$\int_{\mathbf{R}^3} (1, v, |v|^2) f(t, v) dv = \int_{\mathbf{R}^3} (1, v, |v|^2) f_{in}(t, v) dv$$
$$\int_{\mathbf{R}^3} f(t, v) \ln f(t, v) dv \leq \int_{\mathbf{R}^3} f_{in}(v) \ln f_{in}(v) dv$$

for a.e. $t \geq 0$, and

$$\int_{\mathbf{R}^3} f_{in}(v) \phi(0, v) dv + \int_0^T \int_{\mathbf{R}^3} f(t, v) \partial_t \phi(t, v) dv$$
$$= \int_0^T \int_{\mathbf{R}^6} (\Phi(t, v) - \Phi(t, w)) \cdot \nabla(v-w) (F(\nabla_v - \nabla_w) F)(t, v, w) dv dw$$

$$\text{with } \Phi(t, v) := \nabla_v \phi(t, v), \quad F(t, v, w) := \sqrt{\frac{f(t, v) f(t, w)}{8\pi|v-w|}}$$

Notation $\|g\|_{L_k^p}^p := \int (1 + |v|^2)^{k/2} |g(v)|^p dv$ with $p \geq 1$ and $k \in \mathbf{R}$

Definition

A (\mathcal{N}, q, C'_E) -suitable solution on $[0, T) \times \mathbf{R}^3$ is an H-solution s.t.

$$\begin{aligned} & H_+(f(t_2, \cdot) | \kappa) + C'_E \int_{t_1}^{t_2} \left\| \mathbf{1}_{f(t,v) > \kappa} \nabla_v f(t, v)^{1/q} \right\|_{L^q(\mathbf{R}^3)}^2 dt \\ & \leq H_+(f(t_1, \cdot) | \kappa) + 2\kappa \int_{t_1}^{t_2} \int_{\mathbf{R}^3} (f(t, v) - \kappa)_+ dv dt \end{aligned}$$

for all $t_1 < t_2 \in [0, T) \setminus \mathcal{N}$ and $\kappa \geq 1$, where

$$H_+(g | \kappa) := \int_{\mathbf{R}^3} \kappa h_+ \left(\frac{g(v)}{\kappa} \right) dv, \quad h_+(z) := z(\ln z)_+ - (z-1)_+$$

Partial Regularity in Time

Definition A regular time of f , suitable solution on $I \subset (0, +\infty)$, is a time $\tau \in I$ s.t. $f \in L^\infty((\tau - \epsilon, \tau) \times \mathbf{R}^3)$ for some $\epsilon \in (0, \tau)$.
The set of singular (i.e. nonregular) times of f on I is denoted $\mathbf{S}[f, I]$.

Main Thm Let f be a suitable solution to the Landau equation on $[0, T) \times \mathbf{R}^3$ for all $T > 0$, with initial data f_{in} satisfying

$$\int_{\mathbf{R}^3} (1 + |v|^k + |\ln f_{in}(v)|) f_{in}(v) dv < \infty \quad \text{for all } k > 3$$

Then

$$\text{Hausdorff dim } \mathbf{S}[f, (0, +\infty)] \leq \frac{1}{2}$$

Existence Theory

Prop 1 For all $0 \leq f_{in} \in L^1(\mathbb{R}^3)$ s.t.

$$\int_{\mathbb{R}^3} (1 + |v|^k + |\ln f_{in}(v)|) f_{in}(v) dv < \infty \quad \text{for some } k > 3$$

there exists an (\mathcal{N}, q, C'_E) -suitable solution f on $[0, T]$ with initial data f_{in} and

$$C'_E \equiv C'_E[T, q, f_{in}] > 0, \quad q := \frac{2k}{k+3}$$

Desvillettes Theorem

Notation $\|g\|_{L_k^p(\mathbf{R}^N)} := \left(\int_{\mathbf{R}^N} (1 + |v|^2)^{k/2} |g(v)|^p \right)^{1/p}$

Thm For each $0 \leq f \in L_2^1(\mathbf{R}^3)$ s.t. $f \ln f \in L^1(\mathbf{R}^3)$

$$\int_{\mathbf{R}^3} \frac{|\nabla \sqrt{f(v)}|^2 dv}{(1+|v|^2)^{3/2}} \leq C_D + C_D \int_{\mathbf{R}^6} \frac{|\Pi(v-w)(\nabla_v - \nabla_w) \sqrt{f(v)f(w)}|^2}{|v-w|} dv dw$$

with

$$C_D \equiv C_D \left[\int_{\mathbf{R}^3} (1, v, |v|^2, |\ln f(v)|) f(v) dv \right] > 0$$

Corollary Let $0 \leq f_{in} \in L_k^1(\mathbf{R}^3)$ with $k > 2$ s.t. $f_{in} |\ln f_{in}| \in L^1(\mathbf{R}^3)$.

f H-solution s.t. $f|_{t=0} = f_{in} \implies f \in L^\infty(0, T; L_k^1(\mathbf{R}^3))$

(Formal) Truncated H Theorem

One has

$$\begin{aligned} & \frac{d}{dt} H_+(f(t, \cdot) | \kappa) \\ & + \underbrace{\frac{f(t, v)f(t, w)}{16\pi|v-w|} \left| \nabla(v-w) \left(\frac{\mathbf{1}_{f(t,v)>\kappa} \nabla_v f(t, v)}{f(t, v)} - \frac{\mathbf{1}_{f(t,w)>\kappa} \nabla_w f(t, w)}{f(t, w)} \right) \right|^2 dv dw }_{D_1} \\ & = - \int f(t, v)f(t, w) a(v-w) : \nabla_v (\ln \frac{f(t, v)}{\kappa})_+ \otimes \nabla_w (\ln \frac{f(t, w)}{\kappa})_- dv dw \\ & = - \int a(v-w) : \nabla_v f(t, v) \mathbf{1}_{f(t,v) \geq \kappa} \otimes \nabla_w f(t, w) \mathbf{1}_{f(t,w) < \kappa} dv dw \\ & = \int \underbrace{-\operatorname{div}_v(\operatorname{div}_w a(v-w))}_{\geq 0 \text{ (in fact } = \delta(v-w))} (f(t, v) - \kappa)_+ (\kappa - (f(t, w) - \kappa)_-) dv dw \\ & \leq \underbrace{\kappa \int (f(t, v) - \kappa)_+ dv}_{\text{depleted NL}} \end{aligned}$$

Sketch of the Proof of Prop 1

- Replace a with its truncated variant

$$a_n(z) = \frac{1}{8\pi} \left(\frac{1}{|z|} \wedge n \right) \Pi(z), \quad \text{satisfying } \operatorname{div}(\operatorname{div} a_n) \geq 0$$

- Use the Desvillettes theorem to bound

$$\frac{1}{C_D''} \int_{\mathbf{R}^3} \frac{|\nabla_v \sqrt{f(t,v)}|^2}{(1+|v|)^3} \mathbf{1}_{f(t,v)>\kappa} dv \leq D_1 + \int_{\mathbf{R}^3} (f(t,w) - \kappa)_+ dw$$

- Using the Desvillettes corollary with $p' = 2/q$ (recall $q \in (1, 2)$)

$$\begin{aligned} & \left\| \mathbf{1}_{f(t,v)>\kappa} \nabla_v f(t,v)^{1/q} \right\|_{L^q(\mathbf{R}^3)}^q \\ & \leq \left(\frac{2}{q} \right)^q \|f(t, \cdot)\|_{L_{3p/2p'}^p(\mathbf{R}^3)} \left(\int_{\mathbf{R}^3} \frac{|\nabla_v \sqrt{f(t,v)}|^2 \mathbf{1}_{f(t,v)\geq\kappa}}{(1+|v|^2)^{3/2}} dv \right)^{1/p'} \end{aligned}$$

The 1st De Giorgi Type Lemma

Prop 2 Let f be a (\mathcal{N}, q, C'_E) -suitable solution to the Landau equation for $t \in [0, 1]$ with $C'_E > 0$ and $q \in (\frac{6}{5}, 2)$

Then there exists $\eta_0 \equiv \eta_0[q, C'_E] > 0$ s.t.

$$\int_{1/8}^1 H_+(f(t, \cdot)|\tfrac{1}{2}) dt < \eta_0 \implies f(t, v) \leq 2 \quad \text{a.e. on } [\tfrac{1}{2}, 1] \times \mathbf{R}^3$$

Proof of Prop 2

Set

$$\begin{cases} t^k := \frac{1}{2} - \frac{1}{4} \cdot 2^{-k}, & \kappa_k := (1 + (2^{1/q} - 1)(1 - 2^{-k}))^q \\ f_k^+(t, v) := \mu((f(t, v)^{1/q} - \kappa_k^{1/q})_+) & \text{with } \mu(r) := \min(r, r^2) \end{cases}$$

The quantity

$$A_k := \operatorname{ess\,sup}_{t^k \leq t \leq 1} \frac{c_h}{2} \int_{\mathbb{R}^3} f_k^+(t, v)^q dv + \frac{1}{4} C'_E \int_{t^k}^1 \|\nabla_v f_k^+(t, v)\|_{L^q(\mathbb{R}^3)}^2 dt$$

satisfies

$$A_{k+1} \leq M[q, C'_E] \Lambda^k A_k^\beta, \quad \beta := \frac{8}{3} - \frac{2}{q} > 1 \text{ and } \Lambda := 2 \cdot 4^{\frac{5q}{3}}$$

$$\text{so that } A_0 < M^{-\frac{1}{\beta-1}} \Lambda^{-\frac{1}{(\beta-1)^2}} \implies A_k \rightarrow 0 \text{ as } k \rightarrow +\infty$$

- Control A_0 by truncated entropy + conclude by Fatou's lemma

The Improved De Giorgi Type Lemma

Prop 3 Let f be a (\mathcal{N}, q, C'_E) -suitable solution to the Landau equation on $[0, 1]$ with $q \in (\frac{4}{3}, 2)$. Let $\gamma := \frac{5q-6}{2q-2}$. Then there exists $\eta_1 \equiv \eta_1[q, C'_E] > 0$ and $\delta_1 \in (0, 1)$ such that

$$\overline{\lim}_{\epsilon \rightarrow 0^+} \epsilon^{\gamma-3} \int_{1-\epsilon^\gamma}^1 \left\| \mathbf{1}_{f(T,V) > \epsilon^{-\gamma}} \nabla_V f(T, V)^{\frac{1}{q}} \right\|_{L^q(\mathbb{R}^3)}^2 dT < \eta_1$$
$$\implies f \in L^\infty((1 - \delta_1, 1) \times \mathbb{R}^3)$$

- Prop 3 and a Vitali type covering argument imply that

$$\mathcal{H}^{\frac{3-\gamma}{\gamma} = \frac{q}{5q-6}}(\mathbf{S}[f, [\frac{1}{2}, 1]]) < \infty$$

The Main Thm follows by translation+letting $q \rightarrow 2^-$

Proof of Prop 3: (a) Scaling

- Set

$$f_n(t, v) := \epsilon_n^\gamma f(1 + \epsilon_n^\gamma(t - 1), \epsilon_n^\gamma v) \quad \text{with } \epsilon_n := 2^{-n}$$

$$F_n(t, v) := \mu((f_n(t, v)^{1/q} - 1)_+), \quad \int F_n(t, v) dv \leq \epsilon_n^{\gamma-3}$$

- Observe that f_n is a suitable solution of the Landau eq. on $[0, 1]$ with the same constant C'_E as f
- There exists N large enough so that

$$\begin{aligned} n \geq N &\implies \int_0^1 \left(\int |\nabla_v F_n(t, v)|^q dv \right)^{2/q} dt \\ &\leq 4\epsilon_n^{\gamma-3} \int_{1-\epsilon_n^\gamma}^1 \left(\int |\mathbf{1}_{f \geq \epsilon_n^{-\gamma}} \nabla_v f(T, V)^{1/q}|^q dV \right)^{2/q} dT < 8\eta_1 \end{aligned}$$

Proof of Prop 3: (b) Iteration

- Use the Hölder inequality + Sobolev inequality as in the proof of Prop 2, isolating the term $\|\nabla_v F_{n+1}\|_{L_t^2 L_v^q} = O(\eta_1)$ shows that

$$X_m := \operatorname{ess\,sup}_{\frac{1}{2} < t < 1} \int F_{N+m}(t, v)^q dv$$

satisfies

$$X_{m+1} < \rho (\max(1, X_m)^\alpha + \max(1, X_{m-1})^\alpha), \quad X_0, X_1 \leq M$$

$$\text{with } \alpha := q/3, \quad \rho := D(q)\eta_1^{q/2}, \quad M := 2^{(N+3)(3-\gamma)}$$

- With η_1 small so that $\rho < \frac{1}{2}$, an easy induction shows that

$$X_{2m}, X_{2m+1} \leq \max \left(2\rho, (2\rho)^{\frac{1-\alpha^m}{1-\alpha}} M^{\alpha^m} \right) \implies X_{m_0} < 2D(q)\eta_1^{\frac{q}{2}} \ll 1$$

- Hence f_{N+m_0+3} satisfies the assumption in Prop 2, q.e.d.

- The Desvillettes theorem puts the Landau equation in the same class as 3d Navier-Stokes in terms of Lebesgue exponents — except for the $(1 + |v|)^{-3}$ weight

Navier-Stokes	$u \in L_t^\infty L_x^2, \quad \nabla_x u \in L_t^2 L_x^2$
Landau	$\sqrt{f} \in L_t^\infty L_v^2, \quad \nabla_v \sqrt{f} \cap L_t^2 L_{-3}^2$

- Partial regularity in (t, v) à la Caffarelli-Kohn-Nirenberg (1982)?
- Conditional regularity à la Serrin (1962)? for instance

$$\sqrt{f} \in L_t^p L_k^q \text{ with } \frac{2}{p} + \frac{3}{q} < 1 \implies f \in L_{loc}^\infty ?$$

Yes for $p = \infty$ and $k > 5$ [Silvestre 2017, Gualdani-Guillen 2016 for radial nonincreasing solns, Ben Porath 2021 (local in (t, v))]